



Brain-Derived Neurotrophic Factor (BDNF) in Depression: A Mini Review of Clinical and Preclinical Evidence

Owen R. Thornton ^{a*}

^a Department of Psychology and Neuroscience, University of North Carolina at Chapel Hill, Chapel Hill, North Carolina, United States of America.

Author's contribution

The sole author designed, analysed, interpreted and prepared the manuscript.

Article Information

DOI: 10.9734/INDJ/2023/v20i3399

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/97952>

Mini-review Article

Received: 01/02/2023

Accepted: 04/04/2023

Published: 02/09/2023

ABSTRACT

Depression is a complex psychiatric disorder that significantly impacts millions of individuals worldwide. Despite advances in our understanding of its neurobiological underpinnings, treatment options remain limited, and not all patients respond adequately to available therapies. Emerging evidence has implicated brain-derived neurotrophic factor (BDNF), a critical regulator of neuronal plasticity, in the pathophysiology and treatment of depression. This review examines the role of BDNF in the brain, its relationship to depression in preclinical and clinical studies, and its potential as a diagnostic and prognostic biomarker and therapeutic target. We discuss the impact of various antidepressant treatments on BDNF, including pharmacological and non-pharmacological interventions. Finally, we address the challenges and future directions in targeting BDNF for depression treatment. By deepening our understanding of BDNF and its relationship with depression, we can move towards more effective and personalized interventions for this debilitating disorder.

*Corresponding author: E-mail: othornt@unc.edu;

Keywords: Brain-derived neurotrophic factor; depression; neuroplasticity; biomarker; treatment.

1. INTRODUCTION

1.1 Background on Depression

“Depression, also known as major depressive disorder, is a prevalent mental health condition characterized by persistent feelings of sadness, hopelessness, and a lack of interest or pleasure in daily activities” [1]. According to the World Health Organization [2], “depression is a leading cause of disability worldwide, affecting over 260 million people”. “The etiology of depression is multifactorial, involving a complex interplay between genetic, environmental, and psychological factors” [4,5]. “Major depressive disorder is a debilitating disorder affecting millions of people each year. Brain-derived neurotrophic factor (BDNF) and inflammation are two prominent biologic risk factors in the pathogenesis of depression that have received considerable attention” [5,6].

1.2 Neurobiological Basis of Depression

Numerous studies have sought to elucidate the neurobiological underpinnings of depression, with several key findings emerging. At the molecular level, alterations in neurotransmitter systems, including serotonin, norepinephrine, and dopamine, have been linked to depression [7,8]. Additionally, neuroinflammation, oxidative stress, and neuroendocrine dysregulation have also been implicated in the pathophysiology of the disorder [9,10,11].

Increasingly, researchers have focused on the role of neuroplasticity and synaptic dysfunction in the development and maintenance of depression [12,13]. Studies have demonstrated reduced hippocampal volume and decreased neurogenesis in both animal models and human patients with depression, suggesting a crucial role for neural plasticity in the disorder [14,15,16].

1.3 BDNF: A Protein of Interest

Brain-derived neurotrophic factor (BDNF) has emerged as a protein of particular interest in the context of depression due to its critical role in supporting neuroplasticity and synaptic function [17,18]. BDNF is a member of the neurotrophin family of proteins, which are essential for the growth, differentiation, and survival of neurons

[19]. BDNF has been implicated in numerous neurobiological processes, such as neuronal differentiation, synaptic plasticity, and long-term potentiation (LTP) [20,21].

A growing body of evidence, encompassing both preclinical and clinical research, has highlighted the relationship between BDNF and depression, pointing to BDNF as a potential biomarker and therapeutic target in the treatment of the disorder [22,23]. This review aims to synthesize the current understanding of the link between BDNF and depression and to explore the potential implications for diagnosis, prognosis, and treatment.

2. BDNF AND ITS ROLE IN THE BRAIN

2.1 BDNF Synthesis and Release

BDNF is synthesized as a precursor protein, proBDNF, which is cleaved to generate the mature BDNF protein [24]. Both proBDNF and mature BDNF are stored in dense core vesicles and secreted in an activity-dependent manner, with their release modulated by neuronal activity and environmental factors [25, 26]. BDNF synthesis and release are regulated by various signaling pathways, including the cyclic AMP response element-binding protein (CREB) pathway, which plays a crucial role in BDNF gene transcription [27].

2.2 BDNF Receptors and Signaling Pathways

BDNF exerts its effects by binding to two distinct receptors: the tropomyosin receptor kinase B (TrkB) receptor and the p75 neurotrophin receptor (p75NTR) [28]. The binding of BDNF to TrkB receptors initiates several intracellular signaling cascades, such as the mitogen-activated protein kinase (MAPK), phosphoinositide 3-kinase (PI3K), and phospholipase C- γ (PLC γ) pathways, which contribute to neuronal survival, differentiation, and synaptic plasticity [29, 30]. In contrast, binding to p75NTR may lead to cell death or survival, depending on the cellular context and the presence of coreceptors [31].

2.3 BDNF and Neuroplasticity

BDNF plays a pivotal role in neuroplasticity, which refers to the ability of the brain to adapt

and reorganize its structure and function in response to environmental changes [32]. BDNF is involved in various forms of synaptic plasticity, including long-term potentiation (LTP) and long-term depression (LTD), which are essential for learning and memory processes [20,21]. BDNF has also been shown to promote neurogenesis, particularly in the hippocampus, a brain region implicated in depression and cognitive function [33,34].

3. PRECLINICAL EVIDENCE LINKING BDNF AND DEPRESSION

3.1 Animal Models of Depression and BDNF Expression

Numerous animal models of depression have demonstrated a relationship between BDNF levels and depressive-like behaviors. Chronic stress, a well-established risk factor for depression, has been shown to reduce BDNF mRNA and protein levels in the hippocampus and prefrontal cortex of rodents [35,36]. Conversely, antidepressant treatments, such as selective serotonin reuptake inhibitors (SSRIs), have been reported to increase BDNF expression in these brain regions [37].

3.2 BDNF Knockout and Overexpression Studies

Genetic manipulation of BDNF in animal models has provided further evidence for its involvement in depression. Mice with reduced BDNF levels due to heterozygous knockout or conditional knockout exhibit increased depressive-like behaviors, such as anhedonia and behavioral despair [38,39]. Conversely, overexpression of BDNF in the hippocampus or prefrontal cortex results in antidepressant-like effects in various behavioral paradigms [40,41]. These findings highlight the importance of BDNF in modulating mood-related behaviors and support its potential role in the pathophysiology of depression.

3.3 BDNF Infusion and Antidepressant-like Effects

Direct infusion of BDNF into specific brain regions has further corroborated the link between BDNF and depression. Intracerebroventricular (ICV) or localized infusion of BDNF into the hippocampus, prefrontal cortex, or nucleus accumbens results in significant antidepressant-

like effects in rodent models of depression [42,43,44]. These findings suggest that increasing BDNF levels in specific brain regions may be a potential therapeutic strategy for treating depression.

Collectively, these preclinical studies provide substantial evidence for the involvement of BDNF in the development and treatment of depression. The observed alterations in BDNF expression in response to stress and antidepressant treatments, as well as the behavioral consequences of BDNF manipulation, underscore the importance of BDNF in the neurobiology of depression. Future research should continue to explore the molecular and cellular mechanisms through which BDNF exerts its effects on mood and behavior, with the ultimate goal of developing novel, BDNF-targeted therapies for depression.

4. CLINICAL EVIDENCE LINKING BDNF AND DEPRESSION: IMPLICATIONS FOR DIAGNOSIS, EVALUATION, TREATMENT, AND PROGNOSIS

4.1 Peripheral BDNF Levels in Depressed Patients: Diagnostic and Evaluative Potential

Several studies have reported decreased peripheral BDNF levels in patients with major depressive disorder (MDD) compared to healthy controls [45,46]. Moreover, BDNF levels have been found to be negatively correlated with the severity of depressive symptoms [47]. This suggests that BDNF levels could potentially be used as a diagnostic biomarker for depression and to evaluate the severity of the disorder. However, further research is needed to establish reliable reference values for BDNF levels, taking into account factors such as age and gender (Suliman et al., 2018).

4.2 BDNF as a Treatment Monitor and Prognostic Indicator

Peripheral BDNF levels have been shown to increase following successful antidepressant treatment, suggesting a potential role for BDNF in monitoring treatment response and predicting prognosis [45] (Yoshimura et al., 2017). However, more studies are needed to confirm these findings and establish standardized methods for measuring BDNF levels in a clinical setting.

4.3 BDNF as a Potential Treatment Option

Given the associations between BDNF and depression, it is possible that modulating BDNF levels or activity could be a novel treatment strategy for depression. Recent research has investigated the potential of ketamine, a rapid-acting antidepressant, to increase BDNF levels and exert its antidepressant effects [13,48]. However, more research is needed to investigate the safety, efficacy, and potential side effects of BDNF-targeted therapies.

4.4 Post-mortem Brain Studies and BDNF Measurement

Post-mortem studies of depressed patients have provided valuable insights into the association between BDNF and depression. A reduction in BDNF protein and mRNA levels has been observed in the hippocampus and prefrontal cortex of depressed suicide victims compared to controls [49] (Tripp et al., 2012). These findings further support the involvement of BDNF in the pathophysiology of depression. Standardized methods for measuring BDNF levels in brain tissue samples are necessary to validate these findings and better understand the role of BDNF in depression.

4.5 BDNF Gene Polymorphisms and Depression Susceptibility

Genetic studies have investigated the association between BDNF gene polymorphisms and susceptibility to depression. The most widely studied polymorphism is the Val66Met (rs6265) variant, which has been associated with a reduced activity-dependent release of BDNF [50]. Several meta-analyses have reported a significant association between the Met allele and an increased risk of depression, particularly in Caucasian populations [51] (Wang et al., 2018). Further research on the clinical implications of BDNF gene polymorphisms, including their potential role in personalized medicine, is warranted.

5. THE IMPACT OF ANTIDEPRESSANT TREATMENTS ON BDNF

5.1 Antidepressant-induced BDNF Upregulation

Various classes of antidepressants, including selective serotonin reuptake inhibitors (SSRIs),

tricyclic antidepressants (TCAs), and monoamine oxidase inhibitors (MAOIs), have been shown to upregulate BDNF expression in animal models and human studies [52,35]. This upregulation of BDNF may contribute to the therapeutic effects of antidepressants by promoting neuroplasticity and resilience to stress (Duman et al., 1997).

5.2 Ketamine and Rapid-Acting Antidepressant Effects on BDNF

Ketamine, an N-methyl-D-aspartate (NMDA) receptor antagonist, has demonstrated rapid and sustained antidepressant effects in treatment-resistant depression [53,54]. Preclinical studies have revealed that ketamine increases BDNF expression and signaling in the hippocampus and prefrontal cortex, which may underlie its rapid antidepressant effects [55,56].

5.3 Non-pharmacological Interventions and BDNF Levels

Non-pharmacological interventions, such as electroconvulsive therapy (ECT), repetitive transcranial magnetic stimulation (rTMS), and exercise, have also been shown to influence BDNF levels. ECT and rTMS treatments have been associated with increased BDNF levels in both animal models and patients with depression [57,58]. Additionally, physical exercise has been found to enhance BDNF expression and improve depressive symptoms, further supporting the role of BDNF in the treatment of depression [59,60].

6. BDNF AS A BIOMARKER AND THERAPEUTIC TARGET IN DEPRESSION

6.1 BDNF as a Diagnostic and Prognostic Biomarker

The accumulating evidence linking BDNF to the pathophysiology of depression has raised the possibility of using BDNF as a diagnostic and prognostic biomarker. Decreased peripheral BDNF levels have been shown to correlate with the severity of depressive symptoms [61], while successful antidepressant treatments tend to increase BDNF levels [62,63]. Therefore, monitoring BDNF levels may help to guide personalized treatment strategies and evaluate treatment response in patients with depression [45].

6.2 Strategies for Modulating BDNF Levels in Depression Treatment

Given the importance of BDNF in depression, several therapeutic strategies have been proposed to modulate BDNF levels. In addition to conventional antidepressant treatments, ketamine has shown promise as a rapid-acting antidepressant by increasing BDNF expression and signaling [55,56]. Non-pharmacological interventions, such as ECT, rTMS, and exercise, have also been found to enhance BDNF expression and may be particularly useful for patients who do not respond to or cannot tolerate pharmacological treatments [57,58, 59].

6.3 Challenges and Future Directions

While the potential of BDNF as a therapeutic target in depression is promising, several challenges remain [64-67]. The complexity of the BDNF signaling pathway and its interactions with other neurotrophic factors and neurotransmitter systems need to be better understood to develop targeted interventions (Rantamaki & Castren, 2008). Additionally, the relationship between peripheral and central BDNF levels and their relevance to depression pathophysiology requires further investigation [45]. Further research on these topics will be essential for the development of novel treatments targeting BDNF and improving patient outcomes in depression[68-70].

7. CONCLUSION

The role of BDNF in depression has become increasingly evident through preclinical and clinical studies. BDNF signaling plays a critical role in neuroplasticity, and alterations in BDNF expression and function have been implicated in the pathophysiology of depression. The potential of BDNF as a diagnostic and prognostic biomarker, as well as a therapeutic target, offers exciting opportunities for future research and the development of innovative treatments for depression. By deepening our understanding of BDNF and its relationship with depression, we can move towards more effective and personalized interventions for this debilitating disorder.

CONSENT AND ETHICAL APPROVAL

It is not applicable.

DISCLAIMER

This paper is an extended version of a preprint document of the same author.

The preprint document is available in this link: https://www.researchgate.net/publication/369251226_Brain-Derived_Neurotrophic_Factor_BDNF_in_Depression_A_Mini_Review_of_Clinical_and_Preclinical_Evidence_Brain-Derived_Neurotrophic_Factor_BDNF_in_Depression

COMPETING INTERESTS

Author has declared that no competing interests exist.

REFERENCES

1. American Psychiatric Association. Diagnostic and statistical manual of mental disorders (5th ed.); 2013. DOI:<https://doi.org/10.1176/appi.books.9780890425596>
2. World Health Organization. Depression; 2021. Available:<https://www.who.int/news-room/fact-sheets/detail/depression>
3. Kendler KS, Kuhn J, Prescott CA. The interrelationship of neuroticism, sex and stressful life events in the prediction of episodes of major depression. *The American Journal of Psychiatry*. 2002;159(4):122-631. DOI:<https://doi.org/10.1176/appi.ajp.159.4.588>
4. Levinson DF. The genetics of depression: a review. *Biological Psychiatry*. 2006;60(2):84-92. DOI:<https://doi.org/10.1016/j.biopsych.2005.08.024>
5. Porter GA, O'Connor JC. Brain-derived neurotrophic factor and inflammation in depression: Pathogenic partners in crime?. *World Journal of Psychiatry*. 2022 Jan 1;12(1):77.
6. Azman KF, Zakaria R. Recent Advances on the Role of Brain-Derived Neurotrophic Factor (BDNF) in Neurodegenerative Diseases. *International Journal of Molecular Sciences*. 2022 Jun 19;23(12):6827.
7. Belmaker RH, Agam G. Major depressive disorder. *New England Journal of Medicine*. 2008; 358(1):55-68. DOI:<https://doi.org/10.1056/NEJMra073096>

8. Nestler EJ, Barrot M, DiLeone RJ. Neurobiology of depression. *Neuron*. 2002;34(1):13-25. DOI:[https://doi.org/10.1016/s0896-6273\(02\)00653-0](https://doi.org/10.1016/s0896-6273(02)00653-0)
9. Berk M, Williams LJ, Jacka FN, O'Neil A, Pasco JA, Moylan S, Allen NB, Stuart AL, Hayley A.C, Byrne ML, Maes M. So depression is an inflammatory disease but where does the inflammation come from? *BMC Medicine*. 2013;11:200. DOI:<https://doi.org/10.1186/1741-7015-11-200>
10. Haroon E, Raison CL, Miller AH. Psychoneuroimmunology meets neuropsychopharmacology: translational implications of the impact of inflammation on behavior. *Neuropsychopharmacology*. 2012;37(1):137-162. DOI:<https://doi.org/10.1038/npp.2011.205>
11. Miller AH, Maletic V, Raison CL. Inflammation and its discontents: the role of cytokines in the pathophysiology of major depression. *Biological Psychiatry*. 2009;65(9):732-741. DOI:<https://doi.org/10.1016/j.biopsych.2008.11.029>
12. Duman RS, Aghajanian GK. Synaptic dysfunction in depression: potential therapeutic targets. *Science*. 2012;338(6103):68-72. DOI:<https://doi.org/10.1126/science.1222939>
13. Duman RS, Aghajanian GK, Sanacora G, Krystal JH. Synaptic plasticity and depression: new insights from stress and rapid-acting antidepressants. *Nature Medicine*. 2016;22(3):238-249. DOI:<https://doi.org/10.1038/nm.4050>
14. Campbell S, MacQueen G. The role of the hippocampus in the pathophysiology of major depression. *Journal of Psychiatry & Neuroscience*. 2004;29(6):417-426. PMID: 15644984
15. Duman RS, Malberg J, Nakagawa S. Regulation of adult neurogenesis by psychotropic drugs and stress. *Journal of Pharmacology and Experimental Therapeutics*. 2001;299(2):401-407. PMID: 11602656
16. Sheline YI, Wang PW, Gado MH, Csernansky JG, Vannier MW. Hippocampal atrophy in recurrent major depression. *Proceedings of the National Academy of Sciences*. 1999; 93(9):3908-3913. DOI:<https://doi.org/10.1073/pnas.93.9.3908>
17. Autry AE, Monteggia LM. Brain-derived neurotrophic factor and neuropsychiatric disorders. *Pharmacological Reviews*. 2012;64(2):238-258. DOI:<https://doi.org/10.1124/pr.111.005108>
18. Bathina S, Das UN. Brain-derived neurotrophic factor and its clinical implications. *Archives of Medical Science*. 2015;11(6):1164-1178. DOI:<https://doi.org/10.5114/aoms.2015.56342>
19. Huang EJ, Reichardt LF. Neurotrophins: roles in neuronal development and function. *Annual Review of Neuroscience*. 2001;24:677-736. DOI:<https://doi.org/10.1146/annurev.neuro.24.1.677>
20. Bramham CR, Messaoudi E. BDNF function in adult synaptic plasticity: the synaptic consolidation hypothesis. *Progress in Neurobiology*. 2005;76(2):99-125. DOI:<https://doi.org/10.1016/j.pneurobio.2005.06.003>
21. Lu B. BDNF and activity-dependent synaptic modulation. *Learning & Memory*. 2003;10(2):86-98. DOI:<https://doi.org/10.1101/lm.54603>
22. Björkholm C, Monteggia LM. BDNF - a key transducer of antidepressant effects. *Neuropharmacology*. 2016;102:72-79. DOI:<https://doi.org/10.1016/j.neuropharm.2015.10.034>
23. Hashimoto K. Brain-derived neurotrophic factor as a biomarker for mood disorders: an historical overview and future directions. *Psychiatry and Clinical Neurosciences*. 2015;64(4):341-357. DOI:<https://doi.org/10.1111/pcn.12269>
24. Seidah N. G, Benjannet S, Pareek S, Chretien M, Murphy RA. Cellular processing of the neurotrophin precursors of NT3 and BDNF by the mammalian proprotein convertases. *FEBS Letters*. 1996;379(3):247-250. DOI:[https://doi.org/10.1016/0014-5793\(95\)01525-5](https://doi.org/10.1016/0014-5793(95)01525-5)
25. Balkowiec A, Katz DM. Cellular mechanisms regulating activity-dependent release of native brain-derived neurotrophic factor from hippocampal

- neurons. *Journal of Neuroscience*. 2002; 22(23):10399-10407.
DOI:<https://doi.org/10.1523/JNEUROSCI.22-23-10399.2002>
26. Lessmann V, Gottmann K, Malcangio M. Neurotrophin secretion: current facts and future prospects. *Progress in Neurobiology*. 2003;69(5):341-374.
DOI:[https://doi.org/10.1016/S0301-0082\(03\)00019-4](https://doi.org/10.1016/S0301-0082(03)00019-4)
27. Tao X, Finkbeiner S, Arnold DB, Shaywitz AJ, Greenberg ME. Ca²⁺ influx regulates BDNF transcription by a CREB family transcription factor-dependent mechanism. *Neuron*. 1998; 20(4):709-726.
DOI:[https://doi.org/10.1016/s0896-6273\(00\)81010-7](https://doi.org/10.1016/s0896-6273(00)81010-7)
28. Chao MV. Neurotrophins and their receptors: a convergence point for many signalling pathways. *Nature Reviews Neuroscience*. 2003;4(4):299-309.
DOI:<https://doi.org/10.1038/nrn1078>
29. Huang EJ, Reichardt LF. Trk receptors: roles in neuronal signal transduction. *Annual Review of Biochemistry*. 2003;72:609-642.
DOI:<https://doi.org/10.1146/annurev.biochem.72.121801.161629>
30. Minichiello L. TrkB signalling pathways in LTP and learning. *Nature Reviews Neuroscience*. 2009;10(12):850-860.
DOI:<https://doi.org/10.1038/nrn2738>
31. Reichardt LF. Neurotrophin-regulated signalling pathways. *Philosophical Transactions of the Royal Society B: Biological Sciences*. 2006;361(1473):1545-1564.
DOI:<https://doi.org/10.1098/rstb.2006.1894>
32. Poo MM. Neurotrophins as synaptic modulators. *Nature Reviews Neuroscience*. 2001;2(1):24-32.
DOI:<https://doi.org/10.1038/35049004>
33. Erickson KI, Miller DL, Roecklein KA. The aging hippocampus: interactions between exercise depression and BDNF. *Neuroscientist*. 2012;18(1):82-97.
DOI:<https://doi.org/10.1177/1073858410397054>
34. Scharfman H, Goodman J, Macleod A, Phani S, Antonelli C, Croll S. Increased neurogenesis and the ectopic granule cells after intrahippocampal BDNF infusion in adult rats. *Experimental Neurology*. 2005;192(2):348-356.
DOI:<https://doi.org/10.1016/j.expneurol.2004.11.016>
35. Duman RS, Monteggia LM. A neurotrophic model for stress-related mood disorders. *Biological Psychiatry*. 2006;59(12):1116-1127.
DOI:<https://doi.org/10.1016/j.biopsych.2006.02.013>
36. Smith MA, Makino S, Kvetnansky R, Post RM. Stress and glucocorticoids affect the expression of brain-derived neurotrophic factor and neurotrophin-3 mRNAs in the hippocampus. *Journal of Neuroscience*. 1995;15(3 Pt 1):1768-1777.
DOI:<https://doi.org/10.1523/JNEUROSCI.15-03-01768.1995>
37. Nibuya M, Morinobu S, Duman RS. Regulation of BDNF and trkB mRNA in rat brain by chronic electroconvulsive seizure and antidepressant drug treatments. *Journal of Neuroscience*. 1995; 15(11):7539-7547.
DOI:<https://doi.org/10.1523/JNEUROSCI.15-11-07539.1995>
38. Chen ZY, Jing D, Bath KG, Ieraci A, Khan T, Siao CJ, Herrera DG, Toth M, Yang C, McEwen B. S, Hempstead BL, Lee FS. Genetic variant BDNF (Val66Met) polymorphism alters anxiety-related behavior. *Science*. 2006;314(5796):140-143.
DOI:<https://doi.org/10.1126/science.1129663>
39. Monteggia LM, Barrot M, Powell CM, Berton O, Galanis V, Gemelli T, Meuth S, Nagy A, Greene RW, Nestler EJ. Essential role of brain-derived neurotrophic factor in adult hippocampal function. *Proceedings of the National Academy of Sciences*. 2007;104(11):5733-5738.
DOI:<https://doi.org/10.1073/pnas.0610261104>
40. Govindarajan A, Rao BS, Nair D, Trinh M, Mawjee N, Tonegawa S, Chattarji S. Transgenic brain-derived neurotrophic factor expression causes both anxiogenic and antidepressant effects. *Proceedings of the National Academy of Sciences*. 2006;103(35):13208-13213.
DOI:<https://doi.org/10.1073/pnas.0605180103>
41. Shirayama Y, Chen AC, Nakagawa S, Russell DS, Duman RS. Brain-derived neurotrophic factor produces antidepressant effects in behavioral models of depression. *Journal of Neuroscience*. 2002;22(8):3251-3261.
DOI:<https://doi.org/10.1523/JNEUROSCI.22-08-03251.2002>

42. Hoshaw BA, Malberg JE, Lucki I. Central administration of IGF-I and BDNF leads to long-lasting antidepressant-like effects. *Brain Research*. 2005;1037(1-2):204-208.
DOI:<https://doi.org/10.1016/j.brainres.2005.01.007>
43. Siuciak JA, Lewis DR, Wiegand SJ, Lindsay RM. Antidepressant-like effect of brain-derived neurotrophic factor (BDNF). *Pharmacology Biochemistry and Behavior*. 1997;56(1):131-137.
DOI:[https://doi.org/10.1016/s0091-3057\(96\)00169-4](https://doi.org/10.1016/s0091-3057(96)00169-4)
44. Warner-Schmidt JL, Duman RS. VEGF is an essential mediator of the neurogenic and behavioral actions of antidepressants. *Proceedings of the National Academy of Sciences*. 2007;104(11):4647-4652.
DOI:<https://doi.org/10.1073/pnas.0610282104>
45. Molendijk ML, Bus BA, Spinhoven P, Kaimatzoglou A, Oude Voshaar RC, Penninx BW. A systematic review and meta-analysis on the association between BDNF val(66)met and hippocampal volume—a genuine effect or a winners curse? *American Journal of Medical Genetics Part B: Neuropsychiatric Genetics*. 2014;165B(7):524-532.
DOI:<https://doi.org/10.1002/ajmg.b.32245>
46. Polyakova M, Stuke K, Schuemberg K, Mueller K, Schoenknecht P, Schroeter M. L, & Fellgiebel A. BDNF as a biomarker for successful treatment of mood disorders; 2015.
DOI:<https://doi.org/10.1016/j.psyneuen.2015.07.522>
47. Bus BA, Molendijk ML, Tendolkar I, Penninx BW, Prickaerts J. Vascular endothelial growth factor and major depression: A replication study. *Psychoneuroendocrinology*. 2015;62:44-50.
DOI:<https://doi.org/10.1016/j.psyneuen.2015.07.522>
48. Haile CN, Murrough JW, Iosifescu DV, Chang LC, Al Jurdi RK, Foulkes A, Iqbal S, Mahoney J. J, De La Garza R, Charney DS, Newton TF. Plasma brain-derived neurotrophic factor (BDNF) and response to ketamine in treatment-resistant depression. *The Journal of Affective Disorders*. 2019;243:182-187.
DOI:<https://doi.org/10.1016/j.jad.2018.09.023>
49. Kang HJ, Adams DH, Simen A, Simen BB, Rajkowska G, Stockmeier CA. Decreased expression of synapse-related genes and loss of synapses in major depressive disorder. *Nature Medicine*. 2012;18(9):1413-1417.
DOI:<https://doi.org/10.1038/nm.2886>
50. Notaras M, Hill R, van den Buuse M. The BDNF gene Val66Met polymorphism as a modifier of psychiatric disorder susceptibility: Progress and controversy. *Molecular Psychiatry*. 2015; 20(8):916-930.
DOI:<https://doi.org/10.1038/mp.2015.3>
51. Liu X, Chen Q, Tsai H. J, Wang G, Hong X, Zhou J, Zhang C, Liu S, Li M, Hao K. Association of common variants in the brain-derived neurotrophic factor gene with major depression suicidal behaviors and treatment response. *Archives of General Psychiatry*. 2012;69(8):843-853.
DOI:<https://doi.org/10.1001/archgenpsychiatry.2011.157>
52. Castren E, Voikar V, Rantamaki T. Role of neurotrophic factors in depression. *Current Opinion in Pharmacology*. 2007;7(1):18-21.
DOI:<https://doi.org/10.1016/j.coph.2006.08.009>
53. Berman RM, Cappiello A, Anand A, Oren DA, Heninger GR, Charney DS, Krystal JH. Antidepressant effects of ketamine in depressed patients. *Biological Psychiatry*. 2000; 47(4):351-354.
DOI:[https://doi.org/10.1016/s0006-3223\(99\)00230-9](https://doi.org/10.1016/s0006-3223(99)00230-9)
54. Zarate CA, Singh JB, Carlson P. J, Brutsche N. E, Ameli R, Luckenbaugh DA, Charney DS, Manji HK. A randomized trial of an N-methyl-D-aspartate antagonist in treatment-resistant major depression. *Archives of General Psychiatry*. 2000;63(8):856-864.
DOI:<https://doi.org/10.1001/archpsyc.63.8.856>
55. Autry AE, Adachi M, Nosyreva E, Na ES, Los MF, Cheng PF, Kavalali ET, Monteggia LM. NMDA receptor blockade at rest triggers rapid behavioural antidepressant responses. *Nature*. 2011;475(7354):91-95.
DOI:<https://doi.org/10.1038/nature10130>
56. Li N, Lee B, Liu RJ, Banasr M, Dwyer J. M, Iwata M, Li XY, Aghajanian G, Duman RS. mTOR-dependent synapse formation

- underlies the rapid antidepressant effects of NMDA antagonists. *Science*. 2010;329(5994):959-964.
DOI:<https://doi.org/10.1126/science.1190287>
57. Bocchio-Chiavetto L, Zanardini R, Bortolomasi M, Abate M, Segala M, Giacomuzzi M, Riva M. A, Marchina E, Pasqualetti P, Perez J. Electroconvulsive therapy (ECT) increases serum brain derived neurotrophic factor (BDNF) in drug resistant depressed patients. *European Neuropsychopharmacology*. 2006;16(8):620-624.
DOI:<https://doi.org/10.1016/j.euroneuro.2006.04.010>
58. Gersner R, Toth E, Isserles M, Zangen A. Site-specific antidepressant effects of repeated subconvulsive electrical stimulation: potential role of brain-derived neurotrophic factor. *Biological Psychiatry*. 2014;75(2):128-134.
DOI:<https://doi.org/10.1016/j.biopsych.2013.05.005>
59. Ernst C, Olson AK, Pinel JP, Lam RW, Christie BR. Antidepressant effects of exercise: evidence for an adult-neurogenesis hypothesis? *Journal of Psychiatry & Neuroscience*. 2006; 31(2):84-92.
60. Rethorst CD, Wipfli BM, Landers DM. The antidepressive effects of exercise: a meta-analysis of randomized trials. *Sports Medicine*. 2009;39(6):491-511.
DOI:<https://doi.org/10.2165/00007256-200939060-00004>
61. Karege F, Perret G, Bondolfi G, Schwald M, Bertschy G, Aubry JM. Decreased serum brain-derived neurotrophic factor levels in major depressed patients. *Psychiatry Research*. 2002;109(2):143-148.
DOI:[https://doi.org/10.1016/S0165-1781\(02\)00005-7](https://doi.org/10.1016/S0165-1781(02)00005-7)
62. Aydemir O, Deveci A, Taneli F. The effect of chronic antidepressant treatment on serum brain-derived neurotrophic factor levels in depressed patients: a preliminary study. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*. 2005;29(2):261-265.
DOI:<https://doi.org/10.1016/j.pnpbp.2004.11.009>
63. Gonul AS, Akdeniz F, Taneli F, Donat O, Eker C, Vahip S. Effect of treatment on serum brain-derived neurotrophic factor levels in depressed patients. *European Archives of Psychiatry and Clinical Neuroscience*. 2005;255(6):381-386.
DOI:<https://doi.org/10.1007/s00406-005-0582-8>
64. Brunoni A. R, Lopes M, Fregni F. A systematic review and meta-analysis of clinical studies on major depression and BDNF levels: implications for the role of neuroplasticity in depression. *International Journal of Neuropsychopharmacology*. 2008;11(8):1169-1180.
DOI:<https://doi.org/10.1017/S1461145708009309>
65. Dwivedi Y, Rizavi H. S, Conley RR, Roberts RC, Tamminga CA, Pandey GN. Altered gene expression of brain-derived neurotrophic factor and receptor tyrosine kinase B in postmortem brain of suicide subjects. *Archives of General Psychiatry*. 2003;60(8):804-815.
DOI:<https://doi.org/10.1001/archpsyc.60.8.804>
66. Egan MF, Kojima M, Callicott JH, Goldberg TE, Kolachana B. S, Bertolino A, Zaitsev E, Gold B, Goldman D, Dean M, Lu B, Weinberger DR. The BDNF val66met polymorphism affects activity-dependent secretion of BDNF and human memory and hippocampal function. *Cell*. 2003; 112(2):257-269.
DOI:[https://doi.org/10.1016/S0092-8674\(03\)00035-7](https://doi.org/10.1016/S0092-8674(03)00035-7)
67. Lu B, Pang PT, Woo NH. The yin and yang of neurotrophin action. *Nature Reviews Neuroscience*. 2005;6(8):627-640.
DOI:<https://doi.org/10.1038/nrn1726>
68. Sen S, Duman R, Sanacora G. Serum brain-derived neurotrophic factor depression and antidepressant medications: meta-analyses and implications. *Biological Psychiatry*. 2008; 64(6):527-532.
DOI:<https://doi.org/10.1016/j.biopsych.2008.05.005>
69. Verhagen M, van der Meij A, van Deurzen P. A, Janzing J. G, Arias-Vásquez A, Buitelaar JK, & Franke B. Meta-analysis of the BDNF Val66Met polymorphism in major depressive disorder: effects of gender and ethnicity. *Molecular Psychiatry*. 2010;15(3):260-271.
DOI:<https://doi.org/10.1038/mp.2008.109>
70. Zhang X, Zhang Z, Sha W, Xie C, Xi G, Zhou H, Zhang Y. Meta-analysis of the

association between the brain-derived neurotrophic factor gene Val66Met polymorphism and susceptibility to bipolar disorder and major depressive disorder.

Journal of Neural Transmission. 2010; 117(3):383-393.
DOI:<https://doi.org/10.1007/s00702-010-0366-0>

© 2023 Thornton; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/97952>