



Evaluation of Physical, Anti-nutritional and Optimisation of Milling Characteristics of Quinoa Grains for Sustainable Food Security

Mahendra Kumar ^{a*}, Alpana Singh ^a, Anubha Upadhyay ^a
and R. S. Thakur ^a

^a Department of Food Science and Technology, College of Agriculture, Jawaharlal Nehru Krishi Vishwa Vidyalyaya, Jabalpur, M.P.-482004, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2023/v13i113415

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/109011>

Original Research Article

Received: 01/09/2023

Accepted: 06/11/2023

Published: 08/11/2023

ABSTRACT

Quinoa is a popular pseudo-cereal belonging from Andean region of South America. Post-harvest processing of quinoa plays important role in product development, further processing storage and shelf life extension of grains. The aim of present study was evaluation of physical properties, anti-nutritional factors of quinoa and quinoa grain were milled in order to investigate milling yield, milling loss, broken quinoa and head quinoa as the effect of milling in pearling. Post-harvest processing of quinoa was done in the form of pearling after dehulling process and obtained intermediate products. Pearling process of quinoa was done for a constant period of 5 minutes and different milling proportions were evaluated under different runs of the machine. In this study, thousand seed weight of quinoa grain observed about 2.50 to 2.58g in different runs. True density of quinoa ranged between 990 kg/cm³ (Max.) to 983 kg/cm³ (Min.) whereas bulk density of quinoa was stated

*Corresponding author: E-mail: Mahendra1.cdfst@gmail.com;

between 680 to 691 Kg/cm³. Porosity of quinoa showed between 0.302 to 0.308. Saponin content in quinoa samples varied between 3.02 and 3.05% whereas phytin was found about 0.81 to 0.93%. Obtained results also showed L and b value increased after processing while a value decreased. Results obtained from processing showed that husk percentage found maximum as 19.01% whereas highest broken quinoa proportion was 23.63%. It was also observed that highest milling yield or recovery percentage was 63.58% while maximum milling loss recorded as 40.63%. Furthermore, the influence of pearling on the milling characteristics of quinoa were investigated. The milling fractions can be used to obtain new bread products with improved nutritional profiles.

Keywords: Quinoa; pseudo-cereals; pearling; food security; postharvest processing.

1. INTRODUCTION

Food nutrition plays an important role in improving the health status of individuals. Quinoa is a grain like food crop that provided nutrition and sustenance to Andean indigenous cultures for thousands of years and now plays an increasing role in human diets worldwide. Quinoa has been promoted as an alternative agricultural crop due to its stress tolerant characteristics and marketed as a “super food” for its nutritional qualities. A plethora of research has recently emerged on quinoa chemical constituents and therapeutic properties, depicting the crop as an important resource for functional food development [1].

Quinoa (*Chenopodium quinoa* Willd) is an annual herbaceous plant belonging to Amaranthaceae family, but formerly was under Chenopodiaceae family. It originated in the Pacific slopes of the Andes in South America. The Food and Agriculture Organization of the United Nations had declared 2013 as the International Year of Quinoa, which aimed at promoting its worldwide production, consumption, technological development and biodiversity preservation. Presently, it is cultivated in the world in an area of 126 thousand hectares with an annual production of 103 thousand tons. Bolivia in South America is the biggest producer of quinoa with 46 % of world production followed by Peru with 42 % and USA with 6.3 % [2]. In India, quinoa was cultivated in an area of 440 hectares with an average yield of 1053 tons only [3].

Structurally, quinoa is composed of three main parts: the perisperm, the embryo or germ, and the pericarp or seed hull [4]. The perisperm is the primary starch storage portion, the germ is the lipid storage portion, and finally the hull, also called bran, consists mainly of cellulose and hemicellulose. The physicochemical and functional properties of the main components of

quinoa, starch, fibre, and protein, are widely described in the literature [5-7]. The objective of milling is to obtain intermediate products that can be used subsequently in the manufacture of other products.

Milling is a crucial step after post-harvest of quinoa. The basic objective of a quinoa milling system is to remove the husk and the bran layers, and produce an edible, white quinoa kernel that is sufficiently milled and free of impurities. Milled quinoa is comprised of both head quinoa (HQ) and milled quinoa kernels at least three-quarters of their original length [8]. The weights obtained were recorded after each operation. The weights were used to determine the hulling and milling characteristics [9]. Pearling is a mechanical alternative to the washing with water in order to remove saponins. Pearling is a milling technique in which the bran layers are removed sequentially by friction and abrasion operations, which is increasingly recognized by the milling and baking industry. Some studies found that pearled flour has better processing characteristics compared with whole grain flour by direct milling [10].

The need of the hour is to use quality foods for the ever growing and aging population. The focus now is on functional foods with nutraceutical compounds. Hence, quinoa appears to be the best option that fulfills almost all nutritional requirements as it provides essential nutrients that possess health benefits [11].

2. MATERIALS AND METHODS

Good quality raw quinoa grains were procured from the Thakur ji enterprises, Harda, Madhya Pradesh. The different milling products obtained during the processing of raw quinoa are illustrated in Fig. 1.



Fig. 1. Different products of milling obtained during processing of raw Quinoa

2.1 Physical Properties of Quinoa Grains

For determining thousand kernel weight 100 grains were selected randomly from each sample and weighed by means of digital weighing balance (Shimadzu Corporation, Japan, AY120) with an accuracy of 0.001 g, value obtained was then multiplied by 10 [12]. True density defined as the ratio of mass of seeds to their true volume [13]. It was determined using the toluene (C₇H₈) displacement method. Toluene is mainly used instead of water as it is absorbed by seeds to a lesser extent. It also has a benefit of low surface tension and hence fills even shallow dips in seed [14]. The volume of toluene displaced was found by immersing a weighed quantity of seeds in the measured toluene. Approximately 10 g of seeds

were immersed in 25 ml toluene placed in graduated measuring cylinder. Amount of displaced toluene was recorded through the graduated scale of cylinder. The true density was calculated using the equation as below:

$$\text{True Density} \left(\frac{\text{g}}{\text{mL}} \right) = \frac{\text{Weight of seeds (g)}}{\text{Rise in toluene level (mL)}}$$

The bulk density is the ratio of the mass of the grain to its total volume. It was determined by filling a 10 ml measuring cylinder with grain from a height of 15 cm and then weighing the contents by means of a digital electronic balance (Shimadzu Corporation, Japan, AY120) having an accuracy of ± 0.001 g. Bulk density for each replication was calculated from the following relation [13]:

$$\text{Bulk density} \left(\frac{\text{g}}{\text{mL}} \right) = \frac{\text{Weight of the sample(g)}}{\text{Volume occupied by the sample(mL)}}$$

Porosity (ϵ) of bulk seed which is the ratio of free space between grains to total of bulk grains was computed from the values of true density and bulk density as;

$$\text{Porosity} = \frac{(\text{True density} - \text{Bulk density})}{\text{True density}}$$

2.2 Hunter Colour Values of Raw Quinoa Grain

Colour analysis of quinoa breads was done by using Hunter Lab colorimeter (model SM-3001476 micro sensors). The value of a^* ranged from -100 (redness) to +100 (greenness), the b^* values ranged from -100 (blueness) to +100 (yellowness) while as L^* value indicating the measure of lightness, ranged from 0 (black) to 100 (white).

2.3 Milling Characteristics of Quinoa

2.3.1 Dehusking

One of the most crucial steps in the processing of quinoa is the removal of the husks from the grain, which is also known as quinoa pearling. The husk percentage (HP) was calculated by the following formula:

$$\text{Husk (\%)} = \frac{(\text{Weight of quinoa before pearling (gm)} - \text{Weight of quinoa after pearling (gm)}) \times 100}{\text{Weight of quinoa before pearling (gm)}}$$

2.3.2 Milling yield/ recovery

The amount of bran removed from the brown quinoa kernel is measured by the degree of milling. The degree of milling given impacts on milling recovery of quinoa and it influenced consumer's approval. Beyond how much white quinoa is recovered, milling degree affects the colour and cooking characteristics of quinoa. A milling machine should guarantee consistency in the final product's quality, hence raising the raw material's economic value. Weight of polished quinoa includes head and broken also. The Milling percentage (MP) is calculated by the following formula suggested by [9].

$$\text{Milling (\%)} = \frac{\text{Weight of milled quinoa (g)} \times 100}{\text{Weight of rough quinoa (g)}}$$

2.3.3 Head rice percentage of quinoa

The weight percentage of rough quinoa that is still whole after milling is known as the head quinoa yield. Head quinoa recovery (HQR) is a milling quality characteristic that has a significant impact on quinoa market prices. Percentage of head quinoa (excluding broken) obtained from a sample of whole quinoa.

$$\text{Hulling rice recovery of quinoa (\%)} = \frac{\text{Total weight of head quinoa} \times 100}{\text{Total weight of rough quinoa}}$$

2.3.4 Broken quinoa percentage

Broken quinoa is obtained by subtracting head rice quinoa from total quinoa.

2.3.5 Milling loss

Milling loss is obtained by subtraction of milling yield or recovery from total quinoa. Milling loss formula is given below.

$$\text{Milling loss(\%)} = \text{Husk \%} + \text{Broken Quinoa\%} \text{ or } (100 - \text{milling yield \%})$$

2.4 Anti-nutritional Factors Present in Quinoa

2.4.1 Saponin

Total saponin content of the samples was determined according to the gravimetric method given by [15] with slight modifications.

Flour sample (20 g) was mixed with 200 ml of 20% aqueous ethanol solution in a flask and heated with periodic agitation for 4 h at 55°C in a water bath. The mixture was filtered through

Whatman filter paper and the residue was re-extracted with fresh 200 ml of 20% aqueous ethanol. Both the extracts were combined, concentrated to 40 ml at 40°C using rotary evaporator and transferred to a separating funnel and extracted with 20 ml diethyl ether. Re-extraction was done until the aqueous layer was clear in colour and the saponins were then extracted using 60 ml n-butanol.

The butanoic extract was then washed twice with 10 ml of 5% aqueous sodium chloride and evaporated to dryness in a pre-weighed evaporating dish till constant weight. The saponin content was then calculated as percentage of the original sample using formula as follows;

$$\text{Saponin (\%)} = \frac{(W_2 - W_1) \times 100}{\text{Sample weight}}$$

Where,

W₁ = Weight of evaporating dish

W₂ = Weight of evaporating dish + dried extract

2.4.2 Phytic acid

The phytic acid estimation was based on the principle that phytate is extracted With TCA (trichloroacetic acid) and precipitated as ferric salt. The iron content of the precipitate is determined calorimetrically and phytate phosphorous content is calculated from this value assuming a constant 4Fe: 6P molecular ratio in the precipitate. [16].

$$\text{Phytic acid \%} = \frac{\text{uFe(NO}_3\text{)} \times 3 \times 15}{\text{Sample weight}}$$

3. RESULTS AND DISCUSSION

Physical characteristic of quinoa grain plays important role in product formulation and design

of milling equipment. In this study, thousand seed weight of quinoa grain observed about 2.50 to 2.58g in different runs. True density of quinoa ranged between 990 kg/cm³ (Max.) to 983 kg/cm³ (Min.) whereas bulk density of quinoa was stated between 680 to 691 Kg/cm³. Porosity of quinoa showed between 0.302 to 0.308 (Table 1). The results obtained during study for porosity are in close agreement with those reflected for amaranth and quinoa seeds [17,18]. All data were observed significant different from each other. Thousand seed weight of quinoa grain was ranged between 2.52g and 2.58g which was in close agreement with the range 2.5 to 4.1g as recorded the results recorded by [17]. However, [19] reported thousand kernel weight varied from 0.63 to 0.94 g for various varieties of quinoa seed. Bulk density of quinoa grain was varied from 680Kg/m³ to 691 Kg/m³ which was compliance with the 601.3±0.30 kg/m³ demonstrated by [20] for quinoa seeds.

Table revealed that the saponin content in quinoa samples varied between 3.02 and 3.05% whereas phytin was found about 0.81 to 0.93%. All the recorded data were significant different from each other. The best that we can tell, the majority of research has been conducted on pearled quinoa, since pearling has been shown to improve product acceptability by decreasing the amount of saponins [21]. Phytate also adversely affects the absorption of other nutrients such as amino acids, proteins and starch. These results compare well with literature [22].

3.1 Hunter Colour Values of Raw QUINOA GRain

Colour analysis of quinoa grain is important parameter with respect to understand physical characteristics. Presented L, a and B values of hunter colour values in Table 2.

Table 1. Physical properties and anti-nutritional factor of quinoa grain

Physical Properties	Thousand Seed Weight (g)	True density (kg /m ³)	Bulk density (kg /m ³)	Porosity	Saponin (%)	Phytin (%)
R ₁	2.52	983	680	0.308	3.04	0.81
R ₂	2.54	985	687	0.302	3.05	0.92
R ₃	2.58	990	691	0.302	3.02	0.93
SE(m)	0.032	4.082	3.396	0.073	0.046	0.031
CD	N/A	13.03	10.84	N/A	N/A	0.098

R-Replication, CD-Coefficient of dispersion

Table 2. Hunter colour values of raw and processed quinoa grain

S. No.	Hunter Colour Value					
	L		A		b	
	Raw	Milled	Raw	Milled	Raw	Milled
R ₁	49.25	51.10	7.30	6.50	10.20	12.05
R ₂	49.36	49.02	7.22	6.35	10.40	12.15
R ₃	49.45	50.55	7.42	6.10	10.38	12.08

R-Replication

Table 3. Milling characteristics of different samples of quinoa grain

Run	Milling Time (Min)	Husk percentage (%)	Milling yield/recovery (%)	Broken Quinoa (%)	Milling loss (%)
1.	4.20	15.36	61.90	23.07	38.10
2.	4.25	15.42	60.20	22.35	39.80
3.	4.36	16.26	63.58	22.95	36.42
4.	4.55	18.65	63.12	23.63	36.88
5.	4.46	17.54	62.25	23.45	36.75
6.	4.41	16.89	62.87	21.36	37.13
7.	5.02	19.01	61.23	21.85	38.77
8.	4.15	14.86	59.37	20.39	40.63
9.	4.31	15.67	60.49	20.72	39.51
10.	4.50	17.85	61.77	21.56	38.23
SE(m)		0.029	0.035	0.032	0.108
CD		0.087	0.105	0.095	0.322

Table 2 demonstrated that the hunter colour profile of raw quinoa grain increased after processing. L value ranged from 49.25 to 49.45 for raw quinoa grain while milled grain recorded between 49.02 to 51.10. a value of raw and processed quinoa observed from 7.22 to 7.42 and 6.10 to 6.50 respectively. Obtained results showed that b value of raw quinoa and milled quinoa ranged between 10.20 to 10.40 and 12.05 to 12.15 respectively. Similar results were recorded by [23] for L value of wheat and another observations showed at par results [24].

All the results obtained during the course of study tabulated in terms of husk %, milling yield, broken quinoa and milling loss. Table 3 exhibited that husk percentage ranged between 15.36 to 19.01% and maximum found in 7th run while minimum observed in 1st run.

Milling yield presented in table varied from 59.37 to 63.58% in different runs and maximum showed in 3rd run whereas minimum recorded in 8th run.

It was noticed from the table that broken quinoa recorded maximum as 23.63% in 4th run while

minimum observed as 20.39 % in 8th run of pearling. Milling loss revealed by table that it ranged between 36.42 to 40.63% and maximum found in 8th run while minimum observed in 3rd run.

Husk percentage recorded maximum (19.01%) whereas minimum found as (14.86%). Milling Yield/recovery found highest (63.58%) while lowest recorded as (59.37%). The findings of investigation are supported with the reported ranged value from 68.70 to 73.10% with [25], 72.48 to 80.94 with [26] and 69.96 to 71.38 % [27-30]. Broken Quinoa ranged from 20.39% to 23.63%. The earlier reports in case of paddy for broken percentage were 4.28 to 32.40 percent [23] and 31.43 to 33.66 per cent [30]. Milling loss varied between 36.42 % to 40.63%. Further investigations will be necessary to confirm these data.

Depicted figure below (Fig. 2) showed that processing of quinoa affects its various fraction under different processing conditions. Pearling operation demonstrated that milling condition affects husk percentage, broken quinoa percentage and milling loss.




Fig. 2. Effect of milling on its fractions under different runs of pearling

4. CONCLUSION

Post harvesting operations are important for every crop and plays crucial role in the storage and further product formulations. Pearling is done for good product characteristics as it is desirable in various industries for new product development. Present study concluded that milling characteristics of quinoa could influence by pearling process. Bran is also important constitute of quinoa and used for fibre rich product formulations. Physical properties and anti-nutrient factors of quinoa changed during pearling process. By products obtained from the quinoa polishing is majorly used in poultry industry, dairy industry for new product development and research and development. It could be important that mechanization of underutilised grains need to improve. Post-harvest processing factors improve yields of quinoa and reduces wastes due to losses in processing.

ACKNOWLEDGEMENT

I am grateful to my major guide Dr. Alpana Singh, Minor guide Dr. Anubha Upadhyay and all the faculty members of department of food science and technology, JNKVV, Jabalpur who supporting me directly or indirectly.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Zimmet PZ, Magliano DJ, Herman WH, Shaw JE. Diabetes: A 21st century challenge. *Lancet Diabetes Endocrinol.* 2014;2:56–64.
2. FAOSTAT, 2013, Quinoa area and production in the World. Available:<http://www.fao.org>
3. Adolf VI, Shabala S, Andersen MN, Razzaghi F, Jacobsen SE. Varietal differences of quinoa's tolerance to saline conditions. *Plant Soil.* 2012;357:117–129. DOI:10.1007/s11104-012-1133-7
4. Reguera, María, and Claudia Monika Haros. Structure and composition of kernels. *Pseudocereals: Chemistry and technology.* 2017;28-48.
5. Koziol, MJ. Chemical composition and nutritional evaluation of quinoa (*Chenopodium quinoa* Willd). *Journal of Food Composition Analysis.* 1992;5:35-68.
6. Schoenlechner Regine, et al. Functional properties of gluten-free pasta produced from amaranth, quinoa and buckwheat. *Plant foods for human nutrition.* 2010;65:339-349.
7. Kurek MA, Karp S, Wyrwicz J, Niu Y. Physicochemical properties of dietary fibers extracted from gluten-free sources: quinoa (*Chenopodium quinoa*), amaranth (*Amaranthus caudatus*) and millet (*Panicum miliaceum*). *Food Hydrocolloids.* 2018;85:321-330.

8. USDA. United States standards for rice, revised. Washington, D.C.: Federal Grain Inspection Service; 2009.
9. Ghosh AK, Nanda BB, Govindaswami S, Nayak BB. Influence of nitrogen on the physico-chemical characteristics of rice grain. *Oryza*. 1971;8:87–93.
10. Singh S, Singh N. Effect of debranning on the physico-chemical, cooking, pasting and textural properties of common and durum wheat varieties. *Food Res. Int.* 2010;43(9):2277–2283.
11. James LEA. Quinoa (*Chenopodium quinoa* Willd.): Composition, chemistry, nutritional, and functional properties. *Advances in food and nutrition research*. 2009;58:1-31.
12. Varnamkhasti MG, Mobli H, Jafari A, Keyhani AR, Soltanabadi MH, Rafiee S, Kheiralipour K. Some physical properties of rough rice (*Oryza sativa* L.) grain. *Journal of Cereal Science*. 2008;47(3):496–501.
13. Deshpande SD, Bal S, Ojha TP. Physical properties of soybean. *Journal of Agricultural Engineering Research*. 1993;56(2):89-98.
14. Mohsenin NN. Physical properties of plant and animal materials. Structure, physical characteristics and mechanical properties. Physical properties of plant and animal materials. Vol. 1. Structure, physical characteristics and mechanical properties. 1970;1:1.
15. Mir MA, Parihar K, Tabasum U, Kumari E. Estimation of alkaloid, saponin and flavo-noid content in various extracts of *Crocus sativa*. *Journal of Medicinal Plants Studies*. 2016;4(5):171-174.
16. Wheeler EL, Ferrel RE. A method for phytic acid determination in wheat and wheat fractions. *Cereals and grains association*. 1971;48:313-320
17. Vilche, C, Gely M, Santalla E. Physical properties of quinoa seeds. *Biosystems Engineering*. 2003;86(1):59-65.
18. Ghumman A, Mudgal S, Singh N, Ranjan B, Kaur A, Rana JC. Physicochemical, functional and structural characteristics of grains, flour and protein isolates of Indian quinoa lines. *Food Research International*. 2021;140:109982.
19. Gaikwad, KK, Pawar VS, Shingote AB, Shinde EM. Studies physico-chemical properties of Quinoa (*Chenopodium quinoa* willd.) seed. *Pharma. Innov.* 2021;10:612-645.
20. Abalone R, Cassinera A, Gaston A, Lara MA. Some physical properties of amaranth seeds. *Biosystems Engineering*. 2004;89(1):109-117.
21. Gómez-Caravaca, Ana Maria, Giovanna lafelice, Vito Verardo, Emanuele Marconi, Maria Fiorenza Caboni. Influence of pearling process on phenolic and saponin content in quinoa (*Chenopodium quinoa* Willd). *Food chemistry*. 2014;157:174-178.
22. García-Esteba, Rosa Ma, Eduardo Guerra-Hernández, Belén García-Villanova. Phytic acid content in milled cereal products and breads. *Food research international*. 1999;32(3):217-221.
23. Navarro JL. Effect of wheat pearling process on composition and nutritional profile of flour and its bread making performance. *International journal of food science and technology*. 2022;57:249-257.
24. Anderson Jessica. Whole grain wheat effect of peeling and pearling on chemical composition, taste and colour. Master thesis. Swedish university of agri. Sci... Department of food science. 2011;22.
25. El-Bana MA, Goma RA and Abd El-Sattar AS. Effect of parboiling process on milling quality, physical and chemical properties of two rice varieties. *Menoufia J. Food & Dairy Sci*. 2020;5.
26. Kumari MS, D Vijayalakshmi and Usha R. Physical and Cooking Characteristics of Selected Aromatic Rice Varieties. *Journal of Dairying, Foods and Home Sciences*. 2010;29(3and4):227-231.
27. Farahmandfar R, Farahmandfar E, Varnamkhasti MG, Zarei M. Milled rice quality assessment. *International Journal of Food Engineering*. 2010;6(2): Article 7.
28. Verma DK, Mohan M, Prabhakar PK and Srivastav PP. Physicochemical and cooking characteristics of Azad basmati. *International Food Research Journal*. 2015;22(4):1380-1389.
29. Indrasari SD, Purwaningsih, Jumali, Ardhiyanti SD, Handoko DD, Kusbiantoro B. Physical, milling quality and physicochemical properties of several local glutinous paddy varieties in West Java, Indonesia. *IOP Publishing, Earth and Environmental Science*. 2019;309:012014.

30. Rather TA, Malik MA and Dar AH Physical, milling, cooking, and pasting characteristics of different rice varieties grown in the valley of Kashmir India. Cogent Food & Agriculture. 2016; 2:1178694.

© 2023 Kumar et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/109011>