

Pattern of Blood Donation Practices among Students of a Nigerian University

Ugwu, N. I.^{1*}, Ugwu C. N.², Oti W. J. O.³ and Uneke C. J.⁴

¹Department of Haematology and Immunology, Faculty of Clinical Medicine, College of Health Sciences, Ebonyi State University, Abakaliki, Nigeria.

²Department of Internal Medicine, Federal Teaching Hospital, Abakaliki, Nigeria.

³Department of Industrial Chemistry, Faculty of Sciences, Ebonyi State University, Abakaliki.

⁴Directorate of Research, Innovation and Commercialization (DRIC), Ebonyi State University, Abakaliki, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. Author UNI designed the study, performed the statistical analysis, wrote the protocol and wrote the first draft of the manuscript. Authors UCJ and OWJO managed the analyses of the study. Author UCN managed the literature searches. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IBRR/2019/v9i330102

Editor(s):

(1) Dr. Dharmesh Chandra Sharma, Associate Blood Transfusion Officer (ABTO), Incharge Blood Component and Aphaeresis Unit Blood Bank, Department of Pathology, J. A. Groups of Hospital and G. R. Medical College, India.

Reviewers:

(1) Noha Mohamed Kamel, Suez Canal University, Ismailia, Egypt.

(2) J. Y. Peter, University of Abuja, Nigeria.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/49597>

Original Research Article

Received 02 April 2019

Accepted 19 June 2019

Published 26 June 2019

ABSTRACT

Background: Blood transfusion service is an indispensable component of health care. Despite the increased demand for blood, the supply of safe blood has been inadequate.

Objective: The aim of this study is to determine the pattern of blood donation practices among medical students in Abakaliki.

Methods: This was a cross-sectional study carried out at Ebonyi State University Abakaliki between October 2017 and March 2018. Stratified sampling technique was used to recruit participants from medical students using pre-tested, semi-structured, self-administered questionnaire, to assess their pattern of blood donation practices. Data was analysed using SPSS software, version 20.

Results: One hundred and fifty eight medical students who participated in the study were made

up of 90 (58.4%) males and 68 (41.6%) females. The most prevalent age group was 20 to 25years. One hundred and forty three (93%) of the participants were single. The most common blood group was O Rh positive, followed by A Rh positive then B Rh positive while AB Rh negative was the least. Majority of the participants have never donated blood. The minority who have ever donated blood, donated either once or twice and the donation was made for a family member or friend.

Conclusion: Participants were found to have poor pattern of blood donation practices. Misconception, fear, cultural and religious influences deter people from practicing voluntary non-remunerated blood donation. Conducting awareness and enlightenment programs regularly will help to keep them well informed and motivated to practice voluntary non-remunerated blood donation.

Keywords: Voluntary blood donation; students; medical.

1. INTRODUCTION

The timely availability of safe blood products is essential aspect of medical services. Blood is essential to life, circulates through the body and delivers essential substances like oxygen and nutrients to the body cells. It also transports metabolic waste products away from the body cells. Blood transfusion aims at the safe transfer of blood components from the donor to the recipient. In spite of the relevance of blood donation, the major challenge to the transfusion of blood is meeting the increasing demand for blood and ensuring its constant supply. Shortage of blood is due to an increase in the demand, with fewer voluntary blood donors [1,2].

The importance of blood transfusion in medical practice cannot be overemphasized as millions of lives have been saved since the discovery of ABO blood groups. Despite advances in medical research, ideal substitute for blood is yet to be found. Therefore blood donation by humans is still the only source of blood and blood components. Since all blood components and manufactured blood products originate with blood donors, the safety of blood transfusion begins with careful selection of donors. [3] Accordingly, donors should be managed in a way that ensures high standard of care.

WHO estimates that blood donation by 1% of the population is generally the minimum needed to meet a nation's most basic requirements for blood. [4] Generally, donated blood come from either voluntary non-remunerated donors, commercial donors or family members. Family/replacement donors are usually unaware about conditions that may make them unsuitable to donate blood. Paid donors often lead lifestyles that expose them to the risk of infections that

could be transmitted through their blood and are motivated by monetary gain which make them vulnerable to exploitation. [5] An adequate and reliable supply of safe blood can be assured by a stable base of regular, voluntary, unpaid blood donors as they are motivated by altruism or social responsibility and are rewarded with personal satisfaction and self esteem. One of the biggest challenges to blood safety particularly in Sub-Saharan Africa is accessing safe and adequate quantities of blood and blood products. [6] Communities in Africa face several enduring challenges such as chronic blood shortages, high prevalence of Transfusion-Transmissible Infections (TTIs), lack of national blood transfusion services, problems with recruitment and retention of voluntary non-remunerated blood donors, family replacement and commercial blood donation, and inadequate use of pharmacologic and non-pharmacologic alternatives to allogeneic blood. [7] Addressing these challenges should be a central priority for most blood transfusion services, particularly in Sub-Saharan Africa, to ensure the uninterrupted supply of safe blood and blood products. [8] Unlike developed countries, significant percentage of blood donation in developing countries largely depend on family replacement and paid blood donors. [9] According to Nigeria Federal Ministry of Health survey, 25% and 75% of donated blood in public sector is from commercial and family replacement donors while the reverse is the case in private sector where 75% and 25% were commercial and family replacement donors respectively and voluntary blood donors were negligible in both sectors. [10] In our society, overdependence on family replacement and remunerated donors to meet the increasing demand for blood and blood products poses serious danger to potential recipient [11].

Young people are the hope and future of a safe blood supply in the world as they are healthy and enthusiastic. As majority of them will be pursuing their education, schools and colleges can become a good platform for motivational activities. By virtue of operating in health institutions, medical students are expected to be aware of the scarcity of blood and blood products despite increasing demand, the consequences of blood scarcity on health services and are thus expected to donate as well as encourage voluntary blood donation among the public. The objective of this study was to determine the pattern of blood donation practices and the reasons for not donating blood among medical students.

2. MATERIALS AND METHODS

2.1 Study Design and Area

The study was a cross-sectional study carried out at Ebonyi State University, Abakaliki, South Eastern Nigeria between October 2017 and March 2018.

2.2 Study Population, Sample Size and Sampling Technique

The study population comprised medical students who were undergoing their clinical training. Using the Yaro Yamane formula for finite population, a minimum sample size of 169 was calculated. However, the sample size was increased by 10% giving 185 sample size to take care of attrition. The participants were selected using stratified sampling technique. The participants' level of study formed the basis of each stratum. Sampling frame for the study was 304 (comprising 109, 102 and 93 students in 400, 500 and 600 levels respectively). Proportional allocation was used to select the number of students required to partake in the study from each level. At each level, simple random sampling by ballot method was used to select those who participated in the study.

2.3 Data Collection / Data Instrument

The tool for data collection was a semi-structured self-administered questionnaire. The questionnaire consisted of two sections. Section A comprised the socio-demographic characteristics of the participants while section B comprised questions that sought to assess the practice of voluntary blood donation among the participants. The questionnaires were

administered to the students who gave their consent. Questionnaires with incomplete information were excluded.

2.4 Data Analysis

Data was analysed using Statistical Package for Social Sciences (SPSS) software version 20. Descriptive statistics was used to compute mean and standard deviation for continuous data while the categorical data were expressed in frequency and percentages. Results were presented in tables and chart.

3. RESULTS

3.1 Socio-demographic Characteristics

A total of 158 medical students participated in this study. They comprised of 98 males and 60 females with a ratio of 1.6: 1. A higher proportion of the participants 128 (81%) were within the age group 20- 25 years (Table 1).

3.2 Blood Donation Practices among Medical Students

Majority of the students have never donated blood in the past (Fig. 1). Among those who have ever donated blood, most of them are not regular blood donors as majority of them, 69.6% and 19.6% have donated only once and twice respectively. Blood donation was made mainly for either a family member or a friend. Participants (35.4%) who have ever donated blood were mostly males (Table 2).

Some of the reasons given for not donating blood include lack of information on blood donation and it's importance, don't have enough blood to donate, cultural/religious reasons, misuse of blood in the hospital, as shown in Table 3.

4. DISCUSSION

Blood transfusion is a very crucial component of patient management. It has been life-saving procedure especially in cases of medical emergency and for patients suffering from various medical conditions. Voluntary blood donors are the source of safe blood. With the level of ignorance, misconception and fear regarding blood donation, there is a need for medical students to take the lead in this noble course by practicing regular voluntary blood donation to maintain a regular blood supply.

This study revealed that the most prevalent age group among the participants was 20 to 25 years. This collaborates with previous studies which reported that most blood donors are young people [9,12]. In a developing country like Nigeria, lack of information, poor blood donor recruitment and retention strategy and various misconceptions have resulted in a limited number of voluntary donors. This challenge can be tackled by adopting such recruitment methods that can overcome the misconceptions and also motivate the public to donate at an early age, so

that they become lifelong voluntary donors. According to WHO, blood donation can be started safely from 17 years of age, so it is important that the young generation can be motivated to do this harmless task. [13] Young persons are the most potential blood donors in every society and students constitute a huge proportion of them. These potential donors should therefore be well harnessed by sensitization and continued health education using appropriate behaviour change communication models.

Table 1. Socio-demographic characteristics of medical students who participated in the study

Characteristics	Frequency	Percentage
Sex		
Males	98	62
Females	60	38
Total	158	100
Age (years)		
<20	2	1.3
20 – 25	128	81
26 – 30	23	14.6
>30	5	3.2
Total	158	100
Marital status		
Married	7	4.4
Single	151	95.6
Total	158	100
Religion		
Christianity	158	100
Islam	0	0
Traditional	0	0
Others	0	0
Total	158	100




Fig. 1. Medical students who have ever donated blood

Table 2. Blood donation practices among medical students

Variable	Frequency	Percentage
Number of times donated (n = 56)		
1	39	69.6
2	11	19.6
≥3	6	10.8
Blood donation by gender (n = 56)		
Female	10	17.9
Male	46	82.1
Donated without pay (n = 56)		
Yes	49	87.5
No	7	12.5
Who blood was donated for (n = 56)		
Family member (s)/ friend(s)	41	73.2
Stranger(s)	15	26.8

Table 3. Reasons for donating or not donating blood

Responses	Frequency	Percentage
Reasons for donating blood		
To save lives	147	93
As a reward for benefiting from donated blood	6	3.8
To get free medical check	3	1.9
To get money	2	1.3
Total	158	100
Reasons for not donating blood		
Lack of information on blood donation and it's importance	47	29.7
Don't have enough blood to donate	36	22.8
Fear of post-donation outcomes	15	9.5
Fear of needle	13	8.2
Cultural/religious reasons	12	7.6
Never thought of blood donation	10	6.3
Nobody asked for blood donation	8	5.1
Fear of contacting infections	7	4.4
Fear of discovering diseases	5	3.2
Misuse of blood in the hospital	3	1.9
Fear of sight of blood	2	1.3
Total	158	100

This study found that blood donation practice among the participants was poor as majority of them have never donated blood in the past. This corroborates with findings of previous studies which reported poor blood donation practices among the population studied. Kaoje et al. [14] Wedajo et al. [15], in a study conducted at Sokoto, Nigeria reported that only 25% of the participants had ever donated blood.[14] Similarly, a study among health care workers in Ethiopia reported that only 31.9% of the participants had ever donated blood in their life. [15] The poor blood donation practices could be a reflection of perception of the practice of blood donation in the our society.

Another interesting finding of this study is that majority of the people who have ever donated blood were males, with only a few number of females. This corroborates with findings of previous studies which reported minimal contribution to blood donation among females. [16] Similarly, another study also reported that female donors contribute only a few compared to male donors. [17] This is an affirmation of the WHO report that there are more male donors. [13] The reluctance to donate blood among females is understandable because females usually have a lower packed cell volume at certain times of the months due to menstrual flow. Previous studies have reported that women experience 70% more deferral from donation

than men, because of higher frequencies of anaemia, issues related to pregnancy, breast feeding and adverse reactions. [18] Vasovagal attack and post donation fatigue appear to be more common in females compared to males. [19] Hence, general health of women need to be improved by good nutrition and iron supplementation. Considering the fact more females are venturing into the medical profession, the practice of blood donation should be encouraged among females lending credence to the fact that anybody, irrespective of gender, can save lives.

Moreover, majority of those who have ever donated among the participants were first time blood donors as they have donated only once. Similar findings have been reported by previous studies. [20] This suggests that majority of the donors are not regular blood donors and may have donated blood out of need. It is important to note that after the first donation, the students could also donate again if properly motivated.

Majority of the people who have donated did that either to a family member or a friend. Previous studies have given similar report. [21,22] Studies have shown that there is scarcity of regular voluntary blood donors in Nigeria. [9] Many people only donate when there is a compelling need to donate for a family member or a friend. This is different from what is obtainable from developed countries where blood is readily available due to voluntary blood donation practiced regularly by a higher proportion of their population. Globally, it has been found that 80% of first time donors every year give up the practice of blood donation. [23] Reasons given for not donating blood among respondents who had never donated were similar to other studies. [22,24] This may imply that continuous enlightenment is needed among this studied group and indeed, among the general population to highlight the importance of blood donation, especially voluntary non-remunerated blood donation through various channels of communication.

Some of the reasons given by the participants why people do not donate blood voluntarily include lack of information on blood donation and its importance, do not have enough blood to donate, nobody asked for blood donation, cultural and religious reasons, misuse of blood in the hospital, fear of post-donation outcomes. Our finding is similar to that of previous study which reported that the major reasons for not

donating blood were concern about the sterilization of the equipment followed by unknown fear, the collection facility is far from the place, not having enough time to donate. [25,26] Likewise, other studies found that unknown fear of blood donation, needle prick, misuse or selling of their donated blood by the blood bank are some of the reasons for not donating blood. [27, 28] All these reasons can be overcome by encouraging the students to donate blood and educating them about the importance of donating blood. This can be attained by organising different educational programs at that can shed light on the significance of blood donation. With proper education, these erroneous beliefs and misconceptions will be corrected as evidenced from the results of previous studies. [29] This will lead to better understanding and correct information about voluntary blood donation with improved voluntary blood donation practices. The end result will be improvement in the availability of safe blood and blood products for improved medical care services.

This study also revealed that the major reason for donating blood among the participants was to save lives. Other studies conducted in different parts of the globe also reported that the major motivation for donating blood by the participants was the intention to save lives. [21,30] Other reasons given by few of the participants for donating blood include as a reward for benefiting from donated blood, to have free medical check and to get money. Even though the practice of blood donation has been adjudged to be safe, some people still have many wrong ideas about it as they want to donate due to their personal interest and not voluntarily out of altruism. Such donors have been reported not to be safe as they can conceal important medical information in a bid to achieve their goal irrespective of the consequences [5].

5. CONCLUSION

The practice of regular voluntary blood donation among medical students was poor as most of them have never donated blood in the past. Participants who have ever donated blood are not regular blood donors as majority have donated only once. Blood donation in most cases were made for either a friend or a family member. Most of the donors were males as females were more reluctant to donate blood. Some of the reasons given by most of them for donating blood include to save life. Some were motivated because they want to give rewards for

having benefited from donated blood in the past, others to get free medical check while a few were motivated because they want to get money. Some of the reasons given by the participants for not donating blood include lack of information on blood donation and its importance, don't have enough blood to donate, fear of post-donation outcomes, fear of needle, cultural/religious reasons, never thought of blood donation, nobody asked for blood donation, fear of contacting infections, fear of discovering diseases, misuse of blood in the hospital, fear of sight of blood, among others.

There should be a regularly scheduled awareness creation and enlightenment campaigns to allay the fears and misconception related to blood donation. Information on the benefits of regular voluntary blood donation should be emphasized on a continuous basis to correct the impression that blood donation is a harmful practice.

CONSENT

Informed written consent was obtained from each participant before being included in the study.

ETHICAL APPROVAL

Ethical approval was gotten from the Research and Ethics Committee of Ebonyi State University, Abakaliki, Ebonyi State Nigeria.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Aneke JC, Okocha CE. Blood transfusion safety: Current status and challenges in Nigeria. *Asian Journal of Transfusion Science*. 2017;11(1):1-5.
2. Weiping F, Xiaojuan L. Study on the causes and governance path of blood supply shortage in China Mainland. *Science Journal of Public Health*. 2016; 4(3):229- 234.
3. Shrivastava M, Shah N, Navaid S, Agarwal K, Sharma G. Blood donor selection and deferral pattern as an important tool for blood safety in a tertiary care hospital. *Asian J Transfus Sci*. 2016;10(2):122-126.
4. World Health Organisation. Blood transfusion safety; 2006. Available: https://www.who.int/bloodsafety/en/Blood_Transfusion_Safety.pdf [Accessed 02/12/18]
5. Nwogoh B, Awodu OA, Nosakhare BG. Blood donation in Nigeria: Standard of donated blood. *J Lab Physicians*. 2012; 4(2):94-97.
6. Tagny CT, Mbanya D, Tapko JB, Lefrère J. Blood safety in Sub-Saharan Africa: A multi-factorial problem. *Transfusion*. 2008; 48(6):1256–1261.
7. Erhabor O, Adias TC. The challenges of meeting the blood transfusion requirements in Sub-Saharan Africa: The need for the development of alternatives to allogenic blood. *Journal of Blood Medicine*. 2011;2:7-21.
8. Erhabor O, Adias TC, Mainasara AS. Provision of safe blood transfusion services in a low income setting in west Africa: Case study of Nigeria. *Advances in Medicine and Biology*. 2013;59:1-59.
9. Ugwu OA, Madu AJ, Efobi CC, Ibegunam OG. Pattern of blood donation and characteristics of blood donors in Enugu Southeast Nigeria. *Niger J Clin Pract*. 2018;21:1438-43.
10. Federal Ministry of Health. National baseline data survey on blood transfusion. National Blood Transfusion Service. FMOH, Abuja; 2005.
11. Enosolease ME, Imarengiaye CO, Awodu OA. Donor blood procurement and utilization at the University of Benin Teaching Hospital, Benin City. *African Journal of Reproductive Health*. 2004;8(2): 59-63.
12. Pondei K, Lawani E, Pughikumo C. Blood donor practices at two blood banks in Bayelsa State Nigeria. *Journal of Medicine and Medical Sciences*. 2013;4(9):357–361.
13. World Health Organisation. WHO Blood safety and availability fact sheet; 2012. Available: <http://www.who.int/mediacentre/factsheets/fs279/> [Accessed 17 March, 2018].
14. Kaoje AU, Musa AU, Okafoagu NC, Ibrahim H, Raji MO, Ango UM. Public beliefs, Attitude and Practice of Voluntary Non-remunerated Blood Donation among Resident in Sokoto, Nigeria. *Central African Journal of Public Health*. 2017;3(6): 90–96.
15. Wedajo S, Sitote S, Beley M, Reddy PS. Blood donation practice and associated factors among health care workers in public health facility in Dessie Town, North

- East Ethiopia. *Medico Research Chronicles*. 2016;4(6):486–497.
16. Erhabor O, Isaac Z, Abdulrahman Y, Ndakotsu M, Ikhuenbor DB, Aghedo F, et al. Female gender participation in the blood donation process in resource poor settings: Case study of Sokoto in North Western Nigeria. *Journal of Blood Disorders and Transfusion*. 2013;5(1): 1000176.
 17. Bala SS, Handoo S, Jallu AS. Gender differences in blood donation among donors of Kashmir Valley. *Journal of Dental and Medical Sciences*. 2015;14(2): 116–119.
 18. Newmann B. Blood donor suitability and allogeneic whole blood donation. *Transfusion Medicine Review*. 2001;15: 234–244.
 19. Nilsson SB, Sojka P. The blood donation experience: Perceived physical, psychological and social impact of blood donation on the donor. *Vox Sanguinis* 2003;84:120–128.
 20. Jemberu YA, Esmael A, Ahmed KY. Knowledge, attitude and practice towards blood donation and associated factors among adults in Debre Marks town, Northwest Ethiopia. *BMC Hematology*. 2016;16:23-30.
 21. Salaudeen AG, Odeh E. Knowledge and behavior towards voluntary blood donation among students of a tertiary institution in Nigeria. *Nigerian Journal of Clinical Practice*. 2011;14(3):303-307.
 22. Olaiya MA, Alakija W, Ajala A, Olatunji RO. Knowledge, attitudes, Beliefs and motivations towards blood donation among blood donors in Lagos Nigeria. *Transfusion Medicine*. 2004;14(1):13-15.
 23. Thomson RA, Bethel J, Lo AY, Ownby HE, Nass CC, William AE. Retention of Safe blood donors. The retrovirus epidemiology donor study. *Transfusion*. 1998;38:359-367.
 24. Alam M, Masalmeh D. Knowledge, attitudes and practices regarding blood Donation among the Saudi population. *Saudi Medical Journal*. 2004;25(3): 318-21.
 25. Marantidou O, Loukopoulou L, Zervou E, Martinis G, Egglezou A, Fountoli P, et al. Factors that motivate and hinder blood donation in Greece. *Transfusion Medicine*. 2007;17(6):443–450.
 26. Umeora OJJ, Onuh SO, Umeora MC. Socio-cultural barriers to voluntary blood donation for obstetric use in a rural Nigerian village. *African Journal of Reproductive Health*. 2005;9(3):72-76.
 27. Zanin TZ, Hersey DP, Cone DC, Agrawal P. Tapping into a vital resource: Understanding the motivators and barriers to blood donation in Sub-saharan Africa. *African Journal of Emergency Medicine* 2016;6:70–79.
 28. Mishra SM, Sachdev S, Marwaha N, Avasthi A. Study of knowledge and attitude among college going students toward voluntary blood donation from north India. *Journal of Blood Medicine*. 2016;7:19-26.
 29. Vinokur AD, Merion RM, Couper MP, Jones EG, Dong Y. Educational web-based intervention for high school students to increase knowledge and promote positive attitudes towards organ donation. *Health Education and Behaviour*. 2006; 33(6):773–786.
 30. Glynn SA, Kleinman SH, Schreiber GB, Zuck T, Combs SM, Bethel J, et al. Motivations to donate blood: Demographic comparisons. *Transfusion*. 2002;42:216–225.

© 2019 Ugwu et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<http://www.sdiarticle3.com/review-history/49597>