

## **Characterization of Some AUS Rice Genotypes for Yield Traits**

**Md. Kamrul Islam<sup>1</sup>, Md. Shahidur Rashid Bhuiyan<sup>2</sup>, Md. Golam Robbani<sup>2</sup>,  
Md. Sadiquzzaman Sarker<sup>3</sup>, Md. Shamsuzzoha<sup>4</sup> and Maruf Mostofa<sup>1\*</sup>**

<sup>1</sup>*Institute of Seed Technology, Sher-e-Bangla Agricultural University, Dhaka-1207, Bangladesh.*

<sup>2</sup>*Department of Genetics and Plant Breeding, Faculty of Agriculture, Sher-e-Bangla Agricultural University, Dhaka-1207, Bangladesh.*

<sup>3</sup>*Ministry of Agriculture, Dhaka, Bangladesh.*

<sup>4</sup>*Bangladesh Institute of Research and Training on Applied Nutrition, Dhaka-1207, Bangladesh.*

### **Authors' contributions**

*This work was carried out in collaboration among all authors. Authors MKI, MSRB and MGR designed the study, performed the statistical analysis, wrote the protocol, and wrote the first draft of the manuscript. Authors MSS and MS managed the analyses of the study. Author MM managed the literature searches. All authors read and approved the final manuscript.*

### **Article Information**

DOI: 10.9734/AJAHR/2019/v4i130011

#### Editor(s):

(1) Dr. Ahmed Medhat Mohamed Al-Naggar, Professor of Plant Breeding, Department of Agronomy, Faculty of Agriculture, Cairo University, Egypt.

#### Reviewers:

(1) Venkata Sanyasi Seshendra Kumar Karri, Gandhi Institute of Technology and Management, India.

(2) Osama Ali Mohamed Ali, Menoufia University, Egypt.

(3) Ahmed Mohamed Abdelmoghny, Agricultural Research Center, Egypt.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/48923>

**Original Research Article**

**Received 03 March 2019**

**Accepted 17 May 2019**

**Published 20 June 2019**

### **ABSTRACT**

The investigation was carried out under field conditions to characterize yield contributing traits of eighteen (18) advanced Aus rice lines (F<sub>5</sub>) and three commercial check varieties. The experiment was conducted in a randomized complete block design (RCBD). The field was divided into three blocks; each block was sub-divided into 21 plots (lines) where genotypes were randomly assigned. The experiment was conducted during the period of Transplanting Aus season (April 2015 to August 2015) at the genetics and plant breeding experimental field of Sher-e-Bangla Agricultural University, Bangladesh. All the genotypes were characterized and categorized as per the descriptors developed by Biodiversity International, IRRI and WARDA-2007 for DUS test of inbred rice. All the genotypes were grouped and classified as well as described based on yield contributing characters as per descriptors so that all the observed genotypes containing described characters can be easily evaluated and identified at a glance for further studies.

\*Corresponding author: Email: [marufsau@hotmail.com](mailto:marufsau@hotmail.com);

**Keywords:** Aus rice; agro-morphology; Bangladesh; paddy; yield characterization.

## 1. INTRODUCTION

Rice (*Oryza sativa* L.,  $2n = 24$ ) is one of the most broadly cultivated cereal crops in the world spreading across a wide range of geographical, ecological and climatic regions. It belongs to the family Poaceae and subfamily Oryzoidea is the staple food for more than half of the world's population and occupies almost one-fifth of the total land area covered by cereals. High genotypic and phenotypic diversity exists and about more than 120,000 different accessions including landraces are reported in rice globally as a consequence of various adaptations [1]. Most of the world's rice is cultivated and consumed in Asia which constitutes more than half of the world population. It provides 75% of the calories consumed by more than three billion Asians. Approximately 11% of the world's arable land is under rice cultivation and it ranks next to wheat [2].

Rice belongs to the genus *Oryza* and has two cultivated and twenty-two wild species. The two cultivated species are *Oryza sativa* and *Oryza glaberrima*. *Oryza sativa* is grown all over the world while *Oryza glaberrima* has been cultivated in West Africa for the last ~3500 years [3]. Rice is grown under many different conditions and production systems, but submerged in water is the most common method used worldwide. Rice is the only cereal crop that can grow for long periods in standing water [4]. About 57% of rice is grown on irrigated land, 25% on rainfed lowland, 10% on the uplands, 6% in deep water, and 2% in tidal wetlands [5].

Asia can be considered as 'Rice Basket' of the world, as more than 90 percent of the rice is produced and consumed in Asia. World paddy production area was 163.3 million hectares and production were 749.7 million tons [6]. Bangladesh is the 4<sup>th</sup> largest rice producer in the world with the annual production of 345.81 lac metric tons [7]. During Aus season in Financial Year (FY) 2015-16, total cultivated area, production and yield rate of rice was 10,17,969 hectares, 22,88,642 metric tons and 2.248 metric ton per ha respectively. During T. Aman season in Financial Year (FY) 2015-16, total cultivated area, production and yield rate of rice was 55,90,340 hectares, 1,12,39,943 metric tons and 2.412 ton per ha respectively. During Boro season in Financial Year (FY) 2015-16, total cultivated area, production and yield rate of rice

was 47,72,576 hectares, 1,89,37,581 metric tons and 3.968 ton per ha respectively [8].

Several morphological characters are the major determining factors of rice grain yield. Genetic diversity probably serves as an insurance against crop failure [9]. Landraces and wild species possess the immense potential of most valuable genes which can be effectively utilized in the present-day breeding programs to evolve miracle varieties in rice that possess not only high yield potential and quality but also resistant to biotic and abiotic stresses [10].

The total cultivable land is decreasing at a rate of more than 1% per year due to urbanization. The population growth rate is 2 million per year and if the population increases at this rate, the total population will reach 238 million by 2050 [11]. An increase in total rice production is required to feed this ever-increasing population. Aus, Aman, and Boro rice were recently reported to account for 7%, 38%, and 55%, respectively, of the total rice production in Bangladesh [11]. Now, modern high yielding varieties in Aus season are essential to increase the total rice production of Bangladesh. The high yielding varieties of Aus rice are developed through crossing between Aus rice and Boro rice with the intention to increase the yield of Aus rice having genes from Boro rice without much affecting the days to maturity. Eighteen  $F_4$  lines were selected previously which would be used in the present study. The study was undertaken to characterize the  $F_5$  lines which are the prerequisite to release rice variety in future. It will pave the ways for selection of high yielding and short duration Transplant Aus (T. Aus) rice genotypes from eighteen (18) genotypes. The objective of this study to observe high yielding genotypes with their different yield contributing characteristics of T. Aus rice for advanced trials.

## 2. MATERIALS AND METHODS

### 2.1 Experimental Location

The experiment was conducted at the genetics and plant breeding experimental farm of Sher-e-Bangla Agricultural University (SAU), Dhaka, Bangladesh from April 2015 to August 2015. The location of the site was situated at 23°77' N latitude and 90°38' E longitude. Geographically the experimental field is located at 8.4 m above the mean sea level [12].

## 2.2 Climate and Soil

The experimental site was medium high land belonging of Old Madhupur Tract (Agro Ecological Zone - 28) and the soil series was Tejgaon [13]. The soil of the research field was clay loam in texture having pH around 6.5 and organic carbon content is 0.84%. Physical and chemical composition of the experimental field are presented in Table 1. The experiment area was above flood level and having available irrigation and drainage system. The research area was under the subtropical climate. It is characterized by three distinct seasons, winter season from November to February and the pre-monsoon or summer season from March to April and the monsoon period from May to October [14]. Details of the metrological data at the time of experiment were collected from the Bangladesh Meteorological Department (Climate & Weather Division), Agargaon, Dhaka are presented in Table 2 [15].

## 2.3 Planting Materials (Genotypes)

Eighteen (18) genotypes of F<sub>5</sub> generation plus three commercial check varieties (BR 24, BRRI dhan 48 and BRRI dhan 55) were used as experimental materials in the study. Descriptions of the genotypes are given in Table 3.

## 2.4 Experimental Design and Layout

The research was laid out in a randomized complete block design (RCBD). The field was divided into three blocks; each block was subdivided into 21 plots (lines) where genotypes were randomly assigned. The experimental field size was 27 m x 14 m where 1 m border was maintained surrounding the field and every block. The same size of crop field was used for another nine lines. The experimental field was designed in such a way where row to row distance was 25 cm and plant to plant distance was 25 cm. The 18 genotypes plus three check varieties were distributed to each plot within each block randomly.

## 2.5 Collection of Seed

The seeds of 18 Aus lines were collected from germplasm center of Sher-e-Bangla Agricultural University (SAU), Bangladesh. Three check varieties (BR 24, BRRI dhan 48 and BRRI dhan 55) were collected from Bangladesh Rice Research Institute (BRRI).

## 2.6 Application of Fertilizers

The fertilizers N, P, K, S and B in the form of Urea, Triple Super Phosphate (TSP), Muriate of Potash (MOP), Gypsum, Zinc Sulphate and Boron respectively were applied. The entire amount of TSP, MP, Gypsum, Zinc Sulphate and Boron were applied during final preparation of the field. Urea was applied in three equal installments during ploughing, vegetative stage and before flowering. The dose and method of application of fertilizer are shown in Table 4 [16].

## 2.7 Transplanting of Seedling

The seed of all collected rice genotypes soaked separately for 24 hours in a clothes bag. Soaked seeds were picked out from the water and wrapped with straw and gunny bag to increase the temperature for facilitating germination. After 72 hours seeds were sprouted properly. Sprouted seeds were sown separately in the wet seedbed. Proper care was taken so that there was no infestation of pest and diseases and no damage by birds. Healthy seedlings of 25 days old were transplanted in a separate strip of experimental field. Water level was maintained properly after transplanting.

## 2.8 Intercultural Operation and After Care

After the establishment of seedlings, various intercultural operations (irrigation and drainage, gap filling, weeding, top dressing, plant protection measure) were done as per when needed for better growth and development of the rice seedlings.

## 2.9 Harvesting, Threshing and Cleaning

The rice was harvested depending upon the maturity of the plant. Harvesting was done manually from each plot and bundled separately. Properly tagged and brought to the threshing floor. Enough care was taken for threshing and also cleaning of rice seed. Fresh weight of grain was recorded. The grains were cleaned and weight was adjusted to moisture content 14% through drying.

## 2.10 Methods of Recording of Observations

Yield contributing characteristics were collected from ten randomly selected hills from each replicated plot (Table 5). The plants were

selected from the middle of each plot to avoid border effect and portion of the plot. The mean was estimated. Yield contributing traits were recorded using the descriptors developed by BIOVERSITY INTERNATIONAL, IRRI and

WARDA-2007 [17] (Table 7). In addition to the descriptors, the observed genotypes were classified according to Panse and Sukhatme [18]. The observations for characterization were recorded under field condition.

**Table 1. Physical and chemical characteristics of initial soil (0-15 cm depth) of the experimental site**

Physical composition		Chemical composition	
Soil separates	Percent (%)	Soil characteristics	Analytical data
Sand	36.90	Organic carbon (%)	0.84
Silt	26.40	Total N (kg ha <sup>-1</sup> )	1790.00
Clay	36.66	Total S (ppm)	225.00
Texture class	Clay loam	Total P (ppm)	840.00
		Available N (kg ha <sup>-1</sup> )	54.00
		Available P (kg ha <sup>-1</sup> )	69.00
		Exchangeable K (kg ha <sup>-1</sup> )	89.50
		Available S (ppm)	16.00
		pH (1:2.5 soil to water)	5.55
		CEC	11.23

**Table 2. Monthly average temperature, relative humidity and total rainfall of the experimental site during growth period**

Month	Air temperature (°C)		Relative humidity (%)	Total rainfall (mm)
	Maximum	Minimum		
April, 2015	38.0	28.0	79.80	293
May, 2015	37.5	27.0	80.20	307
June, 2015	37.70	27.80	81.08	315
July, 2015	35.45	26.50	83.43	327
August, 2015	34.50	26.00	85.82	338
September, 2015	34.60	25.80	78.08	251

**Table 3. List of genotypes (G) used in the experiment**

Genotypes	Populations	Source
G1	BR 21× BRR I dhan 29, F <sub>5</sub> , S <sub>7</sub> P <sub>5</sub>	SAU
G2	BR 24× BRR I dhan 28, F <sub>5</sub> , S <sub>10</sub> P <sub>10</sub>	SAU
G3	BR 21× BRR I dhan 29, F <sub>5</sub> , S <sub>6</sub> P <sub>3</sub>	SAU
G4	BR 21× BRR I dhan 29, F <sub>5</sub> , S <sub>7</sub> P <sub>2</sub>	SAU
G5	BR 24× BR 26, F <sub>5</sub> , S <sub>6</sub> P <sub>4</sub>	SAU
G6	BR 21× BRR I dhan 29, F <sub>5</sub> , S <sub>6</sub> P <sub>10</sub>	SAU
G7	BR 21× BRR I dhan 29, F <sub>5</sub> , S <sub>7</sub> P <sub>1</sub>	SAU
G8	BR 21× BRR I dhan 29, F <sub>5</sub> , S <sub>6</sub> P <sub>9</sub>	SAU
G9	BR 21× BRR I dhan 29, F <sub>5</sub> , S <sub>7</sub> P <sub>4</sub>	SAU
G10	BR 24× BRR I dhan 28, F <sub>5</sub> , S <sub>10</sub> P <sub>8</sub>	SAU
G11	BR 21× BRR I dhan 36, F <sub>5</sub> , S <sub>1</sub> P <sub>9</sub>	SAU
G12	BR 21× BRR I dhan 29, F <sub>5</sub> , S <sub>6</sub> P <sub>3</sub> (a)	SAU
G13	BR 24× BRR I dhan 29, F <sub>5</sub> , S <sub>5</sub> P <sub>10</sub>	SAU
G14	BR 21× BRR I dhan 29, F <sub>5</sub> , S <sub>6</sub> P <sub>10</sub>	SAU
G15	BR 21× BRR I dhan 29, F <sub>5</sub> , S <sub>6</sub> P <sub>2</sub>	SAU
G16	BR 21× BRR I dhan 29, F <sub>5</sub> , S <sub>1</sub> P <sub>2</sub>	SAU
G17	BR 21× BRR I dhan 29, F <sub>5</sub> , S <sub>6</sub> P <sub>3</sub> (b)	SAU
G18	BR 21× BRR I dhan 29, F <sub>5</sub> , S <sub>6</sub> P <sub>5</sub>	SAU
G19*	BRR I dhan 48	BRR I
G20*	BRR I dhan 55	BRR I
G21*	BR 24	BRR I

F- Generation no; S-Selection no; P-Plant no


**Table 4. Dose and method of fertilizers application in rice field**

Fertilizers	Dose (kg ha <sup>-1</sup> )	Application (%)		
		Basal	1 <sup>st</sup> installment	2 <sup>nd</sup> installment
Urea	127	33.33	33.33	33.33
TSP	52	100	--	--
MOP	60	100	--	--
Gypsum	0	100	--	--
Boron	0	100	--	--

### 2.10.1 Flag leaf: Attitude of the blade

The angle of attachment between the flag leaf blade and the main panicle axis is the attitude of the blade of the flag leaf. It was just visually observed at anthesis period and classified into the following four groups. Flag leaf attitudes are presented in Fig. 1.

Erect (<30°)-1, Intermediate or Semi-erect (30°-45°)-3, Horizontal (46°-90°)-5 and Reflexed or descending (>90°)-7.



**Fig. 1. Flag leaf attitude**

### 2.10.2 Male sterility

It was observed at anthesis period of rice and grouped as per descriptors.

Absent-1, CMS-3, TGMS-5, PGMS-7 and P(T) GMS-9.

### 2.10.3 Microscopic observation of pollen with I<sub>2</sub>-KI solution

It was observed at anthesis period of rice using a microscope and the rice genotypes were classified into eight groups with codes according to guided descriptors as per follow.

Completely sterile with TA pollen-1, Completely sterile with 80% TA pollen-2, Completely sterile

with 50% TA pollen-3, Sterile (91-99%)-4, Partial sterile (31-70%)-5, Partial fertile (31-70%)-6, Fertile (21-30%)-7 and Fully fertile (0-20%)-8.

### 2.10.4 Stigma exertion

Data was observed at anthesis period using a hand lens or magnifying glass and the rice genotypes were classified into five groups with codes according to guided descriptors as per follow.

No or a few (>5%)-1, Low (5-20%)-3, Medium (21-40%)-5, High (41-60%)-7 and Very high (>61%)-9.

### 2.10.5 Panicle curvature of main axis (i.e. recurrent main axis)

Data was collected at near maturity stage and the rice genotypes were classified into four groups with codes according to guided descriptors as per follow.

Absent or very weak (upright)-1, Weak (semi-upright)-3, Medium (slightly drooping)-5 and Strong (strongly dropping)-7.

### 2.10.6 Panicle: Attitude of branches


The compactness of the panicle was classified according to its mode of branching, the angle of primary branches, and spikelet density by the following groups. Attitude of panicle branches are presented in Fig. 2.

Erect (compact panicle)-1, Semi-erect (semi-compact panicle)-3, Spreading (open panicle)-5, Horizontal-7 and Drooping-9.

### 2.10.7 Panicle: Exertion


The extent to which the panicle is exerted above the flag leaf sheath is known as panicle exertion. Data was collected at near maturity stage and the rice genotypes were classified into five groups with codes according to guided

descriptors as per follow. Panicle exertion is presented in Fig. 3.



**Fig. 2. Attitude of panicle branches**

Enclosed-1, Partly exerted-3, Just exerted-5, Moderately exerted-7 and Well exerted-9.



**Fig. 3. Panicle exertion**

**2.10.8 Leaf Senescence: Penultimate leaves are observed at the time of harvest**

Data was collected at the time of harvest and the rice genotypes were classified into three groups with codes according to guided descriptors as per follows.

Late and slow (2 or more leaves retain green color at maturity)-1, Intermediate-5 and Early and fast (leaves are dead at maturity)-9.

**2.10.9 Decorticated Grain: Shape (length-width ratio of de-hulled grain)**

Data was collected at the time of harvest and the rice genotypes were classified into five groups with codes according to guided descriptors as per follows.

Round (L: W<1.5)-1, Bold (L: W=1.5-2.0)-3, Medium (L: W=2.1-2.5)-5, Medium slender (L: W=2.6-3.0)-7 and Slender (L: W>3.0)-9.

**2.10.10 Stem: Culm diameter (from 5 mother tillers in the lowest internode)**


Culm diameter of the stem was measured in millimeter scale at the lowest internode of the stem during flowering or late reproductive stage by using a digital caliper and categorized as per descriptors.

Small (<5.0 mm)-1, Medium (5.1-6.0 mm)-3, Large (6.1-7.0 mm)-5 and Very Large (>7.0 mm)-7

**2.10.11 Stem length (culm length): Measure from the base of the plants to the neck of the panicles**

Stem length (culm length) was measured in centimeter from the base of the plants to the neck of the panicles after flowering to maturity stage and categorized as per descriptors (Fig. 4).

Very short (<40 cm)-1, Short (41–60 cm)-3, Medium (61–80 cm)-5, Long (81-110 cm)-7 and Very long (>110 cm)-9.



**Fig. 4. Culm length**

**2.10.12 Panicle Length: Measured from the neck to the tip of the panicle of main tillers without awns**

The mean length often randomly selected panicles of main tillers from ten hills was measured from neck to the tip of the panicle of the main tiller without awn in centimeters. Data was collected at 7 days after anthesis or full panicle exertion stage According to their length, the observed rice genotypes were classified into four groups with codes.

Short (<20 cm)-3, Medium (21-25 cm)-5, Long (26-30 cm)-7 and Very long (>30 cm)-9.

### 2.10.13 Panicle: Number of the effective tillers per plant

Effective tillers are the tillers which bear panicle and the total number of tillers were counted from each of the sample plants and the average was taken. Based on this character, all the genotypes were grouped into the following groups.

Few (>6)-3, Medium (6-10)-5 and Many (>10)-7.

### 2.10.14 Time of maturity

The total number of days from the date of sowing until 80% seeds became matured considering each replication was recorded on each individual plot and the genotypes were classified as per the guided descriptors.

Very early (>100 days)-1, Early (101-115 days)-3, Medium (116-135 days)-5, Late (136-150 days)-7 and Very late (>150 days)-9.

### 2.10.15 Grain: Weight of 1000 fully developed grains (adjusted of 12% of moisture)

After threshing and recording the net yield, a random sample of fully grown 1000 seeds were counted and weighed at 12% moisture content to record the test weight. According to test weight, the genotypes were categorized into five different groups as follows.

Very low (<15 gm)-1, Low (16-19 gm)-3, Medium (20-23 gm)-5, High (24-27 gm)-7 and Very high (>27 gm) – 9.

### 2.10.16 Grain: Length (without dehulling)

Grain length was measured in mm and a digital caliper was used for clear visualization. Ten grains from every genotype were measured and the mean value was recorded. The genotypes were classified as per the guided descriptors.

Very short (<6.0 mm)-1, Short (6.1-7.0 mm)-3, Medium (7.1-8.0 mm)-5, Long (8.1-9.0 mm)-7 and Very Long (>9.0 mm)-9.

### 2.10.17 Sterile Lemma Length: Measure at post-harvest stage

Sterile lemma length was measured in mm and a digital caliper was used for clear visualization.

Ten grains from every genotype were measured and the mean value was recorded. The genotypes were classified as per the guided descriptors (Fig 5).

Short (<1.5 mm)-1, Medium (1.5-2.5 mm)-3, Long (2.6-3.0 mm)-5 and Very Long (>3.0 mm)-7.




Fig. 5. Rice grain with sterile lemma

### 2.10.18 Decorticated Grain: Length (After dehulling, before milling)

Decorticated grain length was measured in mm and a digital caliper was used for clear visualization (Tab 5). Ten grains from every genotype were measured and the mean value was recorded. The genotypes were classified as per the guided descriptors.

Short (<5.5 mm)-1, Medium (5.6-6.5 mm)-3, Long (6.6-7.5 mm)-5 and Very Long (>7.5 mm)-7.

### 2.11 Statistical Application

The yield contributing data are presented in tabular form for easier description according to the descriptors developed by BIOVERSITY INTERNATIONAL, IRRI AND WARDA-2007 [17]. The data were arranged as per IBPGR-IRRI formulation with the help of the Microsoft Excel 2016 program.

**Table 5. The description of studied traits**

Traits	1	2	3	4	5	6	7	8	9
Flag leaf	Erect (<300)		Semi-erect (300-450)		Horizontal (460-900)		Reflexed (>900)		
Male sterility	Absent		CMS		TGMS		PGMS		P(T) GMS
Microscopic observation of pollen	TA pollen	80% TA pollen	50% TA pollen	Sterile (91-99%)	Partial sterile (31-70%)	Partial fertile (31-70%)	Fertile (21-30%)	Fully fertile (0-20%)	
Stigma exertion	No or a Few (>5%)		Low (5-20%)		Medium (21-40%)		High (41-60%)		Very high (>61%)
Panicle curvature of main axis	Absent or very weak (upright)		Weak (semi-upright)		Medium (slightly drooping)		Strong (strongly drooping)		
Panicle: attitude of branches	Erect		Semi-erect		Spreading		Horizontal		Drooping
Panicle: exertion	Enclosed		Partly exerted		Just exerted		Moderately exerted		Well exerted
Leaf senescence	Late and slow				Intermediate				Early and fast
Decorticated grain	Round (L: W<1.5)		Bold L: W=1.5-2.0)		Medium (L: W=2.1-2.5)		Medium slender (L: W=2.6-3.0)		Slender (L: W>3.0)
Culm diameter	Small (<5.0 mm)		Medium (5.1-6.0 mm)		Large (6.1-7.0 mm)		Very Larg (>7.0 mm)e		
Culm length	Very short (<40 cm)		Short (41-60 cm)		Medium (61-80 cm)		Long (81-110 cm)		Very long (>110 cm)
Panicle length			Short (<20 cm)		Medium (21-25 cm)		Long (26-30 cm)		Very long (>30 cm)
Effective tillers per plant			Few (>6)		Medium (6-10)		Many (>10)		



<b>Traits</b>	<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>5</b>	<b>6</b>	<b>7</b>	<b>8</b>	<b>9</b>
Time of maturity	Very early (>100 days)		Early (101- 115 days)		Medium (116- 135 days)		Late (136-150 days)		Very late (>150 days)
1000 grain weight	Very low (<15 gm)		Low (16-19 gm)		Medium (20-23 gm)		High (24-27 gm)		Very high (>27 gm)
Grain length	Very short (<6.0 mm)		Short (6.1-7.0 mm)		Medium (7.1-8.0 mm)		Long (8.1-9.0 mm)		Very Long (>9.0 mm)
Sterile lemma length	Short (<1.5 mm)		Medium (1.5-2.5 mm)		Long (2.6-3.0 mm)		Very Long (>3.0 mm)		
Decorticated grain	Short (<5.5 mm)		Medium (5.6-6.5 mm)		Long (6.6-7.5 mm)		Very Long (>7.5 mm)		
Panicle length			Short (<20 cm)		Medium (21-25 cm)		Long (26-30 cm)		Very long (>30 cm)
Effective tillers per plant			Few (>6)		Medium (6-10)		Many (>10)		
Time of maturity	Very early (>100 days)		Early (101- 115 days)		Medium (116- 135 days)		Late (136-150 days)		Very late (>150 days)
1000 grain weight	Very low (<15 gm)		Low (16-19 gm)		Medium (20-23 gm)		High (24-27 gm)		Very high (>27 gm)
Grain length	Very short (<6.0 mm)		Short (6.1-7.0 mm)		Medium (7.1-8.0 mm)		Long (8.1-9.0 mm)		Very Long (>9.0 mm)
Sterile lemma length	Short (<1.5 mm)		Medium (1.5-2.5 mm)		Long (2.6-3.0 mm)		Very Long (>3.0 mm)		
Decorticated grain	Short (<5.5 mm)		Medium (5.6-6.5 mm)		Long (6.6-7.5 mm)		Very Long (>7.5 mm)		

**Table 6. Observed genotypes based on culm diameter, culm length, panicle length, time of maturity and thousand grain weight**

<b>Genotypes</b>	<b>Culm diameter (mm)</b>	<b>Culm length (cm)</b>	<b>Panicle length (cm)</b>	<b>Number of effective tillers per plant</b>	<b>Time of maturity (days)</b>	<b>1000 Grain weight (g)</b>
G1	6.28 e	78.52 d	25.71 d	12.23 b-d	103 j	21.27 d-g
G2	5.95 fg	77.64 d	25.57 e	12.90 b-d	109 d-f	22.57 c-e
G3	6.12 ef	78.08 cd	25.52 f	12.80 b-d	106 h-j	21.92 c-g
G4	6.12 e	78.08 d	22.03 s	11.53 d-f	106 ij	21.92 c-f
G5	7.93 a	92.89 ab	23.58 p	12.77 b-d	127.33 b	19.70 fg
G6	6.89 b-d	75.79 de	25.73 c	14.80 a	105.33 g-j	22.27 c-g
G7	6.66 d	78.35 d	25.44 g	12.13 b-e	105.33 g-j	21.73 c-g
G8	6.32 e	78.35 d	25.85 b	12.20 b-d	103.67 ij	22.20 c-g
G9	6.72 cd	69 e	26.40 a	13.73 ab	106 g-i	21.07 e-g
G10	5.18 i	78.79 d	24.67 i	13.27 a-d	110 c-e	23.83 b-d
G11	5.84 gh	57.43 f	20.21 t	15.03 a	107 f-h	19.67 g
G12	5.86 gh	75.88 de	22.91 q	11.63 c-f	105 h-j	21.23 d-g
G13	5.71 gh	92.50 ab	24.66 i	13.43 a-c	130.33 a	23.20 c-e
G14	6.17 ef	78.66 d	23.90 o	9.83 f	108 e-g	23.03 c-e
G15	6.95 bc	95.81 a	24.45 j	12.77 b-d	130.33 a	26.27 ab
G16	6.26 e	69.15 e	24.30 k	10.27 ef	106.33 f-i	23.67 b-e
G17	7.03 b	99.46 a	24.12 m	11.63 c-f	125 b	27.47 a
G18	5.67 h	78.55 d	25.25 h	9.8 f	106 g-i	23.33 c-e
G19*	6.90 b-d	78.61 d	23.95 n	13.53 ab	112 c	24.03 bc
G20*	6.66 d	73.51 de	24.25 l	13.97 ab	111 cd	24.17 bc
G21*	6.75 cd	85.96 bc	22.43 r	12.39 b-d	108 e-g	21.60 c-g
Highest	7.93	99.46	26.40	15.03	130.33	27.47
Lowest	5.18	57.43	20.21	9.80	103	19.67
Mean	6.38	79.57	24.33	12.51	110.98	22.67
CV (%)	2.37	5.33	0.05	9.05	1.63	7.39
LSD <sub>0.05</sub>	0.2496	6.9998	0.0208	1.8671	2.9735	2.7662
Significance level	**	**	**	**	**	**

**Table 7. Classification of the studied twenty one AUS rice genotypes based on the studied eighteen studied traits**

<b>T</b>	<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>5</b>	<b>6</b>	<b>7</b>	<b>8</b>	<b>9</b>	<b>10</b>	<b>11</b>	<b>12</b>	<b>13</b>	<b>14</b>	<b>15</b>	<b>16</b>	<b>17</b>	<b>18</b>
<b>G</b>																		
G1	1	1	7	7	7	3	7	1	9	5	5	7	7	3	5	7	3	3
G2	3	1	7	3	5	5	9	5	9	3	5	7	7	3	5	7	3	3
G3	3	1	7	5	7	3	7	1	7	5	5	7	7	3	5	7	3	3
G4	3	1	7	3	5	3	5	1	9	7	5	5	7	3	5	7	3	3
G5	1	1	7	3	7	3	9	1	9	7	7	5	7	5	3	7	3	5
G6	1	1	7	1	5	3	5	5	7	5	5	7	7	3	5	5	3	3
G7	1	1	7	1	5	3	9	1	9	5	5	7	7	3	5	5	3	3
G8	1	1	7	5	7	3	5	5	9	5	5	7	7	3	5	7	3	3
G9	1	1	7	1	5	3	3	1	9	5	5	7	7	3	5	7	3	3
G10	1	1	7	7	5	3	9	1	7	3	5	5	7	3	7	7	3	3
G11	1	1	7	7	3	3	9	1	9	3	3	5	7	3	3	7	3	3
G12	1	1	7	3	7	3	5	1	7	3	5	5	7	3	5	5	3	3
G13	1	1	7	7	7	3	5	1	9	3	7	5	7	5	5	7	3	5
G14	1	1	7	1	7	3	7	1	7	5	5	5	5	3	5	7	3	3
G15	1	1	7	5	7	3	9	1	9	5	7	5	7	5	7	7	3	5
G16	3	1	7	1	7	3	5	1	9	5	5	5	7	3	7	7	3	3
G17	1	1	7	5	7	3	9	1	9	7	7	5	7	5	7	9	3	5
G18	3	1	7	1	5	3	5	5	9	3	5	7	5	3	5	7	3	3
G19	1	1	7	5	7	3	9	5	9	5	5	5	7	3	7	7	3	3
G20	1	1	7	5	5	3	7	1	9	5	5	5	7	3	7	9	5	7
G21	5	1	7	7	7	3	9	5	9	5	7	5	7	3	5	7	3	3

### 3. RESULTS AND DISCUSSION

#### 3.1 Flag Leaf: Attitude of the blade

From the study, 15 genotypes (G1, G5, G6, G7, G8, G9, G10, G11, G12, G13, G14, G15, G17, G19 and G20) showed erect type flag leaf, 5 genotypes (G2, G3, G4, G16 and G18) showed intermediate or semi-erect type and rest one genotype (G21) showed horizontal type flag leaf.

#### 3.2 Male Sterility

Male sterility was absent in all genotypes (G1, G2, G3, G4, G5, G6, G7, G8, G9, G10, G11, G12, G13, G14, G15, G16, G17, G18, G19, G20 and G21).

#### 3.3 Microscopic Observation of Pollen With I<sub>2</sub>-KI Solution

From the study, all genotypes (G1, G2, G3, G4, G5, G6, G7, G8, G9, G10, G11, G12, G13, G14, G15, G16, G17, G18, G19, G20 and G21) were fertile (21-30%).

#### 3.4 Stigma Exertion

In this case 6 genotypes (G6, G7, G9, G14, G16 and G18) were no or few types, 4 genotypes (G2, G4, G5 and G12) were low type, 6 genotypes (G3, G8, G15, G17, G19 and G20) were medium type and rest 5 genotypes (G1, G10, G11, G13 and G21) were high type for exertion of stigma.

#### 3.5 Panicle Curvature of Main Axis (i.e. recurrent main axis)

One genotype (G11) was observed weak type, 8 genotypes (G2, G4, G6, G7, G9, G10, G18 and G20) were observed medium type and 12 genotypes (G1, G3, G5, G8, G12, G13, G14, G15, G16, G17, G19 and G21) were observed strong type of panicle curvature of the main axis.

#### 3.6 Panicle: Attitude of branches

Where 20 genotypes (G1, G3, G4, G5, G6, G7, G8, G9, G10, G11, G12, G13, G14, G15, G16, G17, G18, G19, G20 and G21) showed semi-erect type and rest 1 genotype (G2) showed spreading type panicle. Erect typed panicles were not found among the genotypes.

#### 3.7 Panicle: Exertion

One genotype (G9) was observed partly exerted type, 7 genotypes (G4, G6, G8, G12, G13, G16 and G18) were observed just exerted type, 4 genotypes (G1, G3, G14 and G20) were observed moderately exerted type and 9 genotypes (G2, G5, G7, G10, G11, G15, G17, G19 and G21) were observed well-exerted type of the panicle exertion. Enclosed type of panicle exertion was not found.

#### 3.8 Leaf Senescence: Penultimate leaves are observed at the time of harvest

Where 16 genotypes (G1, G3, G4, G5, G7, G9, G10, G11, G12, G13, G14, G15, G16, G17 and G20) showed 2 or more leaves retain green color at maturity level and rest 5 genotypes (G6, G8, G18, G19 and G21) showed intermediate type of leaf senescence. Early and fast (leaves are dead at maturity) type of leaf senescence was not found among the genotypes.

#### 3.9 Decorticated Grain: Shape (Length-width Ratio of De-hulled Grain)

Where 16 genotypes (G1, G2, G4, G5, G7, G8, G9, G11, G13, G15, G16, G17, G18, G19, G20 and G21) showed slender type and rest 5 genotype (G3, G6, G10, G12 and G14) showed medium slender type decorticated grain shape. Round, bold and medium type decorticated grain were not found among the genotypes.

#### 3.10 Stem: Culm Diameter (from 5 Mother Tillers in the Lowest Internode)

Culm diameter of observed genotypes ranged from 7.93 mm to 5.18 mm with a mean value of 6.38 mm. As the guided descriptors where there were no small type genotypes on the other hand, six medium type genotypes (G2, G10, G11, G12, G13 and G18), 13 large type genotypes (G1, G3, G4, G6, G7, G8, G9, G14, G15, G16, G19, G20 and G21) and 2 very large type genotypes (G5 and G17) were found (Table 6).

#### 3.11 Stem Length (culm length): Measure from the Base of the Plants to the Neck of the Panicles

Culm length means the length of a stem from ground level to panicle base. Culm lengths of observed genotypes ranged from 99.46 cm to 57.43 cm with a mean value of 79.57 cm. As the guided descriptors where there were no very short type and very long type genotypes on the

other hand, only one short type genotypes (G11), 15 medium type genotypes (G1, G2, G3, G4, G6, G7, G8, G9, G10, G12, G14, G16, G18, G19 and G20) and 5 long type genotypes (G5, G13, G15, G17 and G21) were found (Table 6).

### **3.12 Panicle Length: Measured from the Neck to the Tip of the Panicle of Main Tillers without Awns**

Panicle length of observed genotypes ranged from 26.40 cm to 20.21 cm with a mean value of 24.33 cm. Data were collected at 7 days after anthesis or full panicle exertion stage. As the guided descriptors where there were no short type and very long type genotypes, on the other hand 13 medium type genotypes (G4, G5, G10, G11, G12, G13, G14, G15, G16, G17, G19, G20 and G21) and 8 long type genotypes (G1, G2, G3, G6, G7, G8, G9 and G18) were found (Table 6).

### **3.13 Panicle: Number of the Effective Tillers Per Plant**

The number of effective tillers per plant of the observed genotypes ranged from 15.03 to 9.80 with a mean value of 12.55. There was no genotype showed few types of effective tillers per plant. On the other hand, only two genotypes (G14 and G18) had medium type of effective tillers per plant and rest genotypes (G1, G2, G3, G4, G5, G6, G7, G8, G9, G10, G11, G12, G13, G15, G16, G17, G19, G20 and G21) showed many types of effective tillers per plant (Table 6).

### **3.14 Time of Maturity**

Time of maturity was calculated as days required from sowing to maturity. Time of maturity of the observed genotypes ranged from 130.33 days to 103 days with a mean value of 110.98 days. There was no genotype showed very early, late and very late maturity of the plant. On the other hand, 17 genotypes (G1, G2, G3, G4, G6, G7, G8, G9, G10, G11, G12, G14, G16, G18, G19, G20 and G21) showed early type maturity of plants and rest genotypes (G5, G13, G15 and G17) showed medium type maturity of plants (Table 6).

### **3.15 Grain: Weight of 1000 Fully Developed Grains (adjusted of 12% of moisture)**

Thousand-grain weight of the observed genotypes ranged from 26.27 gm to 19.67 gm

with a mean value of 22.67 gm. In this situation, there was no genotype showed a very low and very high type of 1000 grain weight. On the other hand, 2 genotypes (G5 and G11) showed a low type of 1000 grain weight, 13 genotypes (G1, G2, G3, G4, G6, G7, G8, G9, G12, G13, G14, G18 and G21) showed medium type of 1000 grain weight and rest genotypes (G10, G15, G16, G17, G19 and G20) showed high type of 1000 grain weight (Table 6).

### **3.16 Grain: Length (without Dehulling)**

Grain length of 21 rice genotypes ranged from 10.63 mm to 7.81 mm with a mean value of 8.49 mm. Two genotypes (G17 and G20) were recorded as very long, 16 genotypes (G1, G2, G3, G4, G5, G8, G9, G10, G11, G13, G14, G15, G16, G18, G19 and G21) as long and rest 3 genotypes (G6, G7 and G12) as medium. No genotypes were found as very short and short type.

### **3.17 Sterile Lemma Length: Measure at Post-harvest Stage**

Sterile lemma length of 21 rice genotypes ranged from 2.66 mm to 1.71 mm with a mean value of 1.89 mm. Twenty genotypes (G1, G2, G3, G4, G5, G6, G7, G8, G9, G10, G11, G12, G13, G14, G15, G16, G17, G18, G19 and G21) were recorded as medium and rest one genotype (G20) as long (Tab 7). No genotypes were found as short and very long type.

### **3.18 Decorticated Grain: Length (After Dehulling, before Milling)**

Decorticated grain length of 21 rice genotypes ranged from 8.19 mm to 6.05 mm with a mean value of 6.43 mm. Fifteen genotypes (G1, G2, G3, G4, G6, G7, G8, G9, G10, G11, G12, G14, G16, G18, G19 and G21) were recorded as medium, four genotypes (G5, G13, G15 and G17) as long and rest one genotype (G20) as very long. No genotypes were found as short type.

## **4. CONCLUSION**

Based on yield contributing characteristics developed by Biodiversity International, IRRRI and WARDA-2007 for DUS test of inbred rice, the rice germplasms were classified. All the genotypes were grouped and classified as well as described based on yield contributing characters as per descriptors so that all the

observed genotypes containing described characters can be easily evaluated and identified at a glance for further studies as a part of variety release in future.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Das B, Sengupta S, Parida S, Roy B, Ghosh M, Prasad M, Ghos TK. Genetic diversity and population structure of rice landraces from Eastern and North Eastern States of India. BMC Genetics. 2013;14(1): 71.  
Available:<http://dx.doi.org/10.1186/1471-2156-14-71>
2. Chakravarthi BK, Naravaneni R. SSR marker based DNA finger printing and diversity study in rice (*Oryza sativa*. L.). African J Biotechnol. 2006;5(9):684-688.
3. IRRI (International Rice Research Institute). Rice Research and Production in the 21st Century. (Gramene Reference ID 8380); 2001.
4. IYR (International Year of Rice). Rice and water: A long and diversified story. (Gramene Reference ID 8372). 2004.
5. Forster B. Evolution and Adaptation of Cereal Crops. Edited by V. L. Chopra and S. Prekash. Enfield, NH, USA: Science Publishers Inc (2003), pp. 295. Experimental Agriculture. Cambridge University Press (CUP). 2003;40(1):141–142.  
Available:<http://dx.doi.org/10.1017/s0014479703281527>
6. FAO (Food and Agricultural Organization). Statistical database; 2016.  
Available:<https://www.fao.org/economic/RMM>
7. USDA (United States Department of Agriculture). Statistical database. 2016.  
Available:<https://www.worldriceproduction.com>
8. BBS (Bangladesh Bureau of Statistics). Agriculture crop cutting. Estimation of Aus, T. Aman and Boro rice 2015-2016. Government of the people's Republic of Bangladesh; 2016.  
Available:<http://www.bbs.gov.bd>.
9. Subba Rao LV, Prasad GSV, Prasada Rao U, Rama Prasad A, Acharyulu TL, Rama Krishna S. Collection, Characterization and evaluation of rice germplasm from Bastar Region. Indian J Plant Genet Resour. 2001;14:222-224.
10. Saxena RK, Chang TT, Sapra RL, Paroda RS. Evaluation studies in indigenous rice (*Oryza sativa* L.) germplasm at IRRI. Philippines. Published in NBPGR manual. 1988:1-3.
11. Shelley IJ, Takahashi-Nosaka M, Kano-Nakata M, Haque MS, Inukai Y. Rice cultivation in Bangladesh: Present scenario, problems, and prospects. J Int Coop Agric Dev. 2016;14(4): 20-29.
12. Mostofa M, Roy TS, Chakraborty R, Modak S, Kundu PK, Zaman MS, Rahman M, Shamsuzzoha M. Effect of vermicompost and tuber size on processing quality of potato during ambient storage condition. Int J Plant Soil Sci. 2019;26(3):1-18.  
Available:<http://dx.doi.org/10.9734/ijpss/2018/46554>
13. UNDP (United Nations Development Programme). Land Resources Appraisal of Bangladesh for Agricultural Development. Report 2. Agroecological Regions of Bangladesh. United Nations Development Programme and Food and Agriculture Organization. 1988;212-221.
14. Edris KM, Islam ATMT, Chowdhury MS, Haque AKMM. Detailed Soil Survey of Bangladesh, Dept. Soil Survey and Govt. People's Republic of Bangladesh. 1979: 118.
15. Meteorological Department, Dhaka, Bangladesh; 2015.  
Accessed 13 September 2016.  
Available: <http://www.bmd.gov.bd/>
16. BRRI (Bangladesh Rice Research Institute). Adhunik Dhaner Chash (In Bangoli). Bangladesh Rice Research Institute, Joydebpur, Gazipur. 2015: 5-10.
17. Biodiversity International, IRRI and WARDA. Descriptors for wild and cultivated rice (*Oryza* spp.). Bioversity International, Rome, Italy; International

Rice Research Institute, Los Banos, Philippines; WARDA, Africa Rice Center, Cotonou, Benin; 2007. 18. Panse VG, Sukhatme PV. Statistical methods for agricultural workers, Rev Edn. ICAR, New Delhi; 1995.

---

© 2019 Islam et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*  
*The peer review history for this paper can be accessed here:*  
<http://www.sdiarticle3.com/review-history/48923>