



Excess Parameters of Binary Mixtures of 1, 2-Dimethoxyethane-Propylene Carbonate and 2-Methoxyethyl Ether-Propylene Carbonate

John N. Obowu¹, Millicent U. Ibezim-Ezeani^{1*} and Augustine A. Abia¹

¹*Department of Pure and Industrial Chemistry, University of Port Harcourt, P.M.B. 5323, Choba, Port Harcourt, Nigeria.*

Authors' contributions

This work was carried out in collaboration between all authors. Author AAA designed the study. Author JNO performed the statistical analysis, wrote the protocol and wrote the first draft of the manuscript and author MUIE managed the analyses of the study and literature reviews. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/PSIJ/2018/42560

Editor(s):

- (1) Dr. Marian Apostol, Professor, Theoretical Physics, Institute for Atomic Physics, Magurele-Bucharest, Romania.
(2) Dr. Roberto Oscar Aquilano, School of Exact Science, National University of Rosario (UNR), Rosario, Physics Institute (IFIR) (CONICET-UNR), Argentina.

Reviewers:

- (1) Harekrushna Sutar, Indira Gandhi Institute of Technology, India.
(2) P. Paul Divakar, Sir C. R. Reddy College, Affiliated to Adikavi Nannaya University, India.
(3) Anil Kumar Koneti, GVR&S College of Engineering & Technology, India.
Complete Peer review History: <http://www.sciencedomain.org/review-history/25509>

Original Research Article

Received 23rd April 2018
Accepted 5th July 2018
Published 11th July 2018

ABSTRACT

The excess parameters associated with liquid mixtures were evaluated in order to ascertain the relative strength of intermolecular interactions in binary mixtures of 1,2-Dimethoxyethane-Propylene Carbonate (DME-PC) and 2-Methoxyethyl ether-Propylene Carbonate (DGM-PC). The density and viscosity of pure components of Propylene Carbonate (PC), 1,2-Dimethoxyethane (DME), 2-Methoxyethyl ether (DGM) and mixtures (DME-PC and DGM-PC) were measured within the composition range of 0 to 100% at temperatures of 298.15, 308.15, 318.15, 328.15 and 338.15 K. Experimental density and viscosity values decreased with an increase in system temperature and mole fractions of DME and DGM. Mathematical operations were conducted from the density and viscosity data at 298.15 K to resolve the magnitude of excess molar volume (V_m^E), excess viscosity (η^E), excess Gibbs free energy of activation of viscous flow (G^{*E}) and Grunberg-Nissan interaction parameter (d). These parameters were mainly of negative value, which signified the

*Corresponding author: E-mail: millicent.ibezim-ezeani@uniport.edu.ng;

dominance of dipole-dipole interactions within DME-PC and DGM-PC solvent systems. The existence of interactions between the components of the binary mixtures was also reflected by the non-zero result findings for values of excess parameters.

Keywords: Excess volume; excess viscosity; excess Gibbs free energy of activation of viscous flow; Grunberg-Nissan parameter; dipole-dipole interaction.

1. INTRODUCTION

Technically, the need to augment:

- (i) ionic dissociation and conductivity
- (ii) solubility and mobility of species
- (iii) durability and thermal stability of cell
- (iv) reduction in free energy of reaction or interaction (polar and non-polar)

are the fundamental reasons for deploying mixed solvents in electrochemical systems. This allows for adjustment of properties of the medium of interest by varying proportions of the components of the mixture. Ionic salts are characterized by large lattice energy and considerable energy expenditure is required to separate the ions when they dissolve, giving solutions comprising of discrete anions and cations dispersed in the solvent. Countless salts dissolve in water and in other polar non-aqueous medium, indicating the favourable nature of interactions between solvent molecules and dissolved ions. The introduction of ions into polar solvents is beneficial from the standpoint of ion but may possibly not be from that of solvent. As a rule, solvents with pronounced intermolecular forces (dipolar interactions and hydrogen bonding) are efficient ion solvates. In the solvation of ions, the solvent structure is largely modified owing to strong commove of solvent-solvent interactions in the vicinity of an ion. The knowledge of thermodynamic properties helps in formulating principles for the molecular design of electrolytes utilized in high energy-density batteries. Denticity of coordinating solvent molecules affect the stability of solvated complexes [1]. The ideal number of ligands results from combination of factors namely: the consequence of the contest between attraction to ionic field, steric repulsion forces between coordinating solvent groups and entropy effects of the ligands associated with the metal ion [2]. Viscosity (transport property) and dielectric constant (bulk property of matter) are indispensable parameters in the formulation of state-of-the-art conductive electrolytes for industrial electrochemical processes [3,4]. Earlier reports stated that the low viscosity property of

solvent supports to lessen resistance for ion transport in the solution, while high dielectric constant assists in the reduction of energy of ionic dissociation [5,6]. In studying thermodynamic properties, the interpretation of excess parameters is obligatory to comprehend intermolecular interactions amid components of the mixture. Intermolecular interactions can be established by the deviation of density, viscosity and volume properties from ideality. Thermodynamic properties of binary mixtures find application in many chemical processing industries. Direct experimental measurements are frequently employed for studying non-ideal mixtures, requiring measurements performed over different composition range. By appraising thermodynamic properties and studying volume properties, we can construe the processes that occur in mixtures. The design of electrochemical and engineering processes comprising mass transfer, fluid flow and heat transfer require critical data provided by the excess molar volumes of component mixtures. New consequences resulting from mixing processes can influence volume behaviour such as structural changes of components of the mixture and intermolecular attraction or repulsion between unlike molecules. Profound insight into transport properties can be achieved by studying viscosity behaviour of liquid mixtures. Viscosity and density data are extensively used in solution theory, equipment modelling and molecular dynamics. In the coating and pharmaceutical industries, formulation processes require the benefit of viscosity data. Excess thermodynamics properties have been employed by researchers for studying the relative potency of intermolecular interaction in liquid mixtures [7]. The interactions at the molecular level and its influence on the viscosity and density measurements in binary solvent systems of DME-PC and DGM-PC at various temperatures and compositions were considered in this case study. The glymes (DME and DGM) are aprotic saturated diethers that exhibit extreme solvency and stability in strong bases and moderate stability in acid solutions. DME and DGM efficiently solvate cations, surging anion reactivity and as such can upturn reaction rates. Their diether structures engender

weak associations between the molecules, and this is responsible for the low viscosity and astounding wetting properties of these solvents. The structural feature of DME and DGM that contributes to their functionality relates to the format of oxygen atoms, as ether linkages at two-carbon intervals. This structural arrangement, analogous to that of crown ethers, gives these solvents the capacity to form complexes with myriad of cations. This study was carried out to investigate the deviation of liquid mixtures from ideality and to construe the relationship between molecular interaction and properties (density and viscosity) of the mixtures. The resultant inference will lay the groundwork for the formulation of electrolyte system with high density chemical source of current using DME-PC and DGM-PC as solvates.

2. MATERIALS AND METHODS

The sourced solvents for this research are DGM (99.5%) product of BASF Chemical Company England, DME (99.5%) and PC (99.5%) products of BDH Chemical Ltd. England and dry Mg (BH₄)₂ (95%) product of Sigma Aldrich. The Binary mixtures of DME & PC and DGM & PC were prepared in varying volume percent proportions of 0, 15, 25, 50, 75, 85 and 100%. These were constituted by mixing appropriate volumes of the solvent components in standard flasks fitted with caps and the mass measurements were performed on HR-120 (A&D Japan) electronic balance with a precision of $\pm 10^{-4}$ g. The mole fractions range from 0.125 to 1.000 (DME) and 0.095 to 1.000 (DGM). The density of the pure solvents and mixtures was determined with a single stem capillary pycnometer bottle at the temperatures 298.15, 308.15, 318.15, 328.15 and 338.15 K. The kinematic viscosity of the pure solvents and mixtures was determined using a suspended ubbelohde viscometer tube ASTM D 446. Koehler electronic constant temperature water bath was used to maintain the experimental solvents and mixtures at the desired temperatures of 298.15, 308.15, 318.15, 328.15 and 338.15 K \pm 0.01 K. The flow time was determined electronically by using an electronic timer with precision of \pm 0.01 s. The kinematic viscosity was calculated using the formula [8]:

$$V = kt \quad (1)$$

where V , k and t are the kinematic viscosity, viscometer constant and flow time respectively.

The kinematic viscosity was converted into dynamic viscosities using the formula:

$$\eta = V\rho \quad (2)$$

where η and ρ are the dynamic viscosity and density of liquid respectively. The process was repeated until three successive determinations were obtained within ± 0.1 seconds.

The equations [9] for excess molar volume (V_m^E), excess viscosity (η^E), excess Gibbs free energy of activation of viscous flow (G^{*E}) and Grunberg-Nissan interaction parameter (d) are as follows:

$$V_m^E = X_1 M_1 (1/\rho_m - 1/\rho_1) + X_2 M_2 (1/\rho_m - 1/\rho_2) \quad (3)$$

$$\eta^E = \eta_m - (X_1 \eta_1 + X_2 \eta_2) \quad (4)$$

where X_1 and X_2 are mole fractions of pure constituents 1 and 2 respectively, M_1 and M_2 are molar masses of pure constituents 1 and 2 respectively, ρ_1 and ρ_2 are densities of pure constituents 1 and 2 respectively, η_1 and η_2 are viscosities of pure constituents 1 and 2 respectively, ρ_m and η_m are density and viscosity of the mixture respectively.

$$V_m = \frac{X_1 M_1 + X_2 M_2}{\rho_m} \quad (5)$$

$$G^{*E} = RT[\ln \eta_m V_m - (X_1 \ln \eta_1 V_1 + X_2 \ln \eta_2 V_2)] \quad (6)$$

where R is the Universal constant for gases, T is Absolute temperature, V_1 , V_2 and V_m are molar volumes of pure constituents 1, 2 and mixture respectively.

$$d = \frac{\ln \eta_m - (X_1 \ln \eta_1 + X_2 \ln \eta_2)}{X_1 X_2} \quad (7)$$

3. RESULTS AND DISCUSSION

3.1 Density

The density of the solvent mixtures (DME-PC and DGM-PC) decreased with decrease in PC content of the mixture (Figs. 1 and 2). DME and DGM are aprotic, polar and self-associated solvents having dipole moments of 1.71 and 1.91 D respectively. It is likely that on mixing PC and DME / DGM, the molecular order in PC decreased due to the polar influence of DME / DGM molecules, while the molecular association in pure DME / DGM was disrupted by

interactions with PC. This disruption of molecular order and association within the binary systems is responsible for the decrease in density of the mixtures. Furthermore, as DME proportion in the mixture increases, intermolecular dipolar forces within PC are disrupted, creating weaker interactions between the molecules of PC and DME / DGM; with the result that molecules drift relatively apart after mixing, leading to reduction in density of the solvent mixture. The density of the binary mixtures (DME-PC and

DGM-PC) decreased as temperature increased from 298.15 to 338.15 K (Figs. 1 and 2). It could be that increase in temperature introduced some level of stress and change on the internal energy of the system. The effect of this external influence on the forces of molecular attraction is the distortion in orientation of the solvent-solvent dipoles; which is reflected in the decline of density at higher system temperatures.




Fig. 1. Plot of Density versus mole fraction (DME)




Fig. 2. Plot of Density versus mole fraction (DGM)

3.2 Viscosity

Increase in the temperature of the binary mixtures (DME-PC and DGM-PC), increased the average speed of molecules, leading to decrease in the amount of time spent in contact with neighbouring molecules. As temperature increased, the average intermolecular forces decreased and thus, accounts for the observed decrease in viscosity of the mixtures (DME-PC and DGM-PC) with temperature increase (Figs. 3

and 4). The viscosity of the binary mixture also decreased as the ratio of DME or DGM in the mixture increased, due to the lower viscosity of the solvents. Furthermore, the measured values of viscosity for DME-PC were lower than those of DGM-PC at all experimental temperatures. This is likely due to the higher steric influence experienced as a result of the structurally longer DGM molecule on interaction with PC as compared to that of the relatively shorter chain length of DME.




Fig. 3. Plot of viscosity versus mole fraction (DME)




Fig. 4. Plot of viscosity versus mole fraction (DGM)

3.3 Excess Parameters

The measure of variation between the values of property of solution thermodynamics and that for an ideal solution under parallel conditions of temperature and composition is termed the excess property. This level of variation represents the positive or negative excess of solution thermodynamics, relative to the ideal solution as reference [10]. Excess parameters have zero value for ideal mixing. The negative or positive values of excess parameters indicate non-ideal behavior in mixtures, signifying the existence of specific interactions between the mixtures of DME-PC and DGM-PC. The positive values of excess volume are attributed to dispersive forces, while negative values are attributed to dipole-dipole interactions and formation of hydrogen bonds [11]. Negative values of excess volume result from strong interactions between unlike molecules (DME-PC and DGM-PC); while positive excess volume results from strong interactions between like molecules (DME-DME, DGM-DGM and PC-PC), indicating breakup interactions between unlike molecules of the mixtures. Positive values of excess volume are the result of rupture of hydrogen bonds and loosening of dipole interactions [12]. The non-zero results for values of the excess parameters in this study point towards the existence of interactions between the components of the binary mixtures (Figs. 5 to 8). Various workers have explained the nature and strength of these interactions on the grounds of the sign and magnitude of excess parameters [13]. In both binary solvent systems studied, one factor which may contribute to excess molar volume is the disruption of the intermolecular dipolar forces in component 2 (PC) by the addition of component 1 (DME or DGM). The values of V_m^E for the mixtures (Fig. 5) are

negative over the entire constitution range and it appears that intermolecular interactions between unlike molecules predominate over intermolecular interactions between like molecules. The molar volumes of PC and DME molecules are 85.22 and 104.74 cm³ mol⁻¹ while that for PC and DGM are 85.22 and 142.44 cm³ mol⁻¹ respectively. It is apparent that the accommodation of one component into the structure of the other interstitially is not favorable enough. Therefore, it may not significantly contribute towards the V_m^E values. Thus, dipolar and dispersion forces between DME-PC and DGM-PC molecules are primarily responsible for negative values of V_m^E [14].

According to Fort and Moore [15], the negative values of η^E suggest the existence of dispersion forces; thus, negative deviation of η^E suggests that dipole interactions are more probable in DME-PC and DGM-PC (Fig. 6) systems.

To understand the forces acting between the unlike molecules of DME-PC and DGM-PC, the support of excess Gibbs free energy of activation of flow (G^{*E}) and Grunberg-Nissan interaction parameter (d) is needed. The negative values of excess Gibbs free energy of activation for viscous flow (Fig. 7) indicate the dominance of dispersion forces characterized by decrease in the internal energy of viscous flow; revealing the incidence of weak interaction in the systems. Positive deviations in values of excess activation energy of flow indicate specific interactions between the components of the mixture [16]. The Gibbs free energy of activation for viscous flow is negative over the constitution range (0.125 to 0.710 mole fraction of DME and 0.095 to 0.772 mole fraction of DGM), indicating the dominance of dispersion forces.




Fig. 5. Plot of molar excess volume versus mole fraction




Fig. 6. Plot of excess viscosity versus mole fraction




Fig. 7. Plot of excess gibbs free energy versus mole fraction




Fig. 8. Plot of Grunberg-Nissan parameter versus mole fraction

The sign and magnitude of Grunberg-Nissan interaction d-parameter is considered the measure of strength of interactions. The negative values of the d-parameter (Fig. 8) show the existence of weak interactions between the unlike molecules DME-PC and DGM-PC. Furthermore, positive values of the d-parameter indicate that interactions between unlike molecules are strong [16]. The resolved negative trend of d-parameter values in most of the experimental cases is suggestive of dipole-dipole and dipole-induced dipole interaction of little magnitude between species of dissimilar molecules (DME-PC and DGM-PC) [9].

4. CONCLUSION

The study on excess parameters of binary mixtures (DME-PC and DGM-PC) has revealed its usefulness in understanding the nature of intermolecular interactions between two liquids. Parameters such as density and viscosity of pure solvents and binary liquid mixtures measured at different temperatures over 0 to 100% composition range provide statistics on thermodynamic properties associated with heat and fluid flow. Dipolar and dispersion forces in DME-PC and DGM-PC molecules are principally responsible for negative values of V_m^E and η^E , and it appears that intermolecular interactions between unlike molecules predominate over that of like molecules. The perusal of the sign and magnitude of G^{*E} and d , indicated that interactions between dissimilar molecules in both binary systems studied to follow the order: DME-PC > DGM-PC. This means that solvent-solvent interactions between unlike molecules are stronger in the DME-PC system than in DGM-PC system. Thus, DGM molecules are relatively free from hindrances credited to strong dispersion and dipolar forces between unlike molecules. DGM molecules will, therefore, exhibit great affinity for the electrolyte, and easily stabilize cations in solution, which in turn affords better conducting electrolytes.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Truong NG, Norris AR, Shin H S, Buncel E, Bannard RAB, Purdon JG. Selectivities and thermodynamic parameters of alkali-metal and alkaline-earth-metal complexes of polyethylene-glycol dimethyl ethers in methanol and acetonitrile. *Inorg. Chim. Acta.* 1991;184(1):59-65.
2. Deetz D, Cao F, Wang Q, Sun H. Exploring the liquid structure and ion formation in magnesium borohydride electrolyte using density functional theory. *J. Electrochem. Soc.* 2018;165(2):61-70.
3. Wang P, Anderko A, Young RD. Modeling viscosity of concentrated and mixed-solvent electrolyte systems. *Fluid Phase Equilibria.* 2004;226:71-82.
4. Logan ER, Tonita EM, Gering KL, Li J, Ma X, Beaulieu LY, Dahn JR. A study of the physical properties of Li-Ion battery electrolytes containing esters. *J. Electrochem. Soc.* 2018;165(2):21-30.
5. Matsuda Y, Nokashima H, Morita M, Takasu Y. Behaviour of some ions in mixed organic electrolytes of high energy density batteries. *Electrochem. Sci. Technol.* 1981;128(12):2552-2556.
6. Ding MS, XU K, Jow TR. Conductivity and viscosity of PC-DEC and PC-EC solutions of LiBOB. *J. Electrochem. Soc.* 2005; 152(1):132-140.
7. Bahadur I, Letcher TM, Singh S, Redhi G, Venkatesu P, Ramjugernath D. Excess molar volumes of binary mixtures (an ionic liquid + water): A review. *J. Chem. Thermodyn.* 2015;82:34-46.
8. Ibezim-Ezeani MU, Menegbo LI, Abia AA. Physic-chemical and mixing parameters of electrolyte system containing 1, 2-dimethoxyethane and ethylene carbonate. *Int. J. Emerging Technol. And Advanced Eng.* 2017;7(2):9-14.
9. Ibezim-Ezeani, MU, Menegbo LI. Thermodynamics of electrolyte system with binary mixture of 1, 2-dimethoxyethane and ethylene carbonate. *Int. J. Current Res.* 2017;9(4):48725-48729.
10. Mesquita FMR, Feitosa FX, de Santiago-Aguiar RS, de Sant'Ana HB. Experimental density data and excess molar volumes of coconut biodiesel + n-hexadecane and coconut biodiesel + diesel at different temperatures. *Braz. J. Chem. Eng.* 2014; 31(2):548-551.
11. Sujata SP, Sunil RM. Thermodynamic study of binary mixtures of methyl acrylate and ethyl acrylate with octane-1-ol. *J. Advances Appl. Sci. Res.* 2013;4(3):329-334.
12. El-Banna M. Excess molar volumes and viscosities of binary-mixtures of propylene carbonate with (C-5-C-8) n-alkanols at

- 298.15 K. Canadian. J. Chem. 1997; 75(12):1890-1895.
13. Sati R, Choudhary SN, Prasad K, Mishra VK. Interaction study of polar + nonpolar liquid system: binary mixtures of MPK + Benzene and MPK + cyclohexane. Process Solid State Physics Symposium, Varanasi, India. 1991;196-197.
14. Prakash KM, Dilip KH. Density and viscosity for propylene carbonate + 1,2-dimethoxyethane at 298.15, 308.15, and 318.15 K. J. Chem. Eng. Data. 1994; 39(2):375-377.
15. Fort RJ, Moore WR. Viscosities of binary liquid mixtures. Trans. Faraday Soc. 1966; 62:1112-1119.
16. Sathyanarayana B, kumar BR, Jyostna TS, Satyanarayana N. Densities and Viscosities of Binary Mixtures of N-methylacetamide with some chloroethanes and chloroethenes at T=308.15k. J. Chemical Thermodyn. 2007;39(1):16-21.

© 2018 Obowu et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

*The peer review history for this paper can be accessed here:
<http://www.sciencedomain.org/review-history/25509>*